

ST7 (h3): 293T Lysate: sc-173958

BACKGROUND

ST7 (suppressor of tumorigenicity 7 protein), also known as HELG, RAY1, SEN4, TSG7, ETS7q or FAM4A1, is a 585 amino acid protein localized to the cell membrane. ST7 is ubiquitously expressed, with highest levels found in liver, heart and pancreas. Expressed as seven isoforms produced by alternative splicing events, ST7 may act as a tumor suppressor. The gene that encodes ST7 maps to human chromosome 7q31.2, which encodes over 1,000 genes and makes up about 5% of the human genome. Chromosome 7 has been linked to osteogenesis imperfecta, Pendred syndrome, lissencephaly, citrullinemia and Shwachman-Diamond syndrome. The deletion of a portion of the q arm of chromosome 7 is associated with Williams-Beuren syndrome, a condition characterized by mild mental retardation, an unusual comfort and friendliness with strangers and an elfin appearance. Deletions of portions of the q arm of chromosome 7 are also seen in a number of myeloid disorders including cases of acute myelogenous leukemia and myelodysplasia.

REFERENCES

1. Zenklusen, J.C., et al. 2001. Mutational and functional analyses reveal that ST7 is a highly conserved tumor-suppressor gene on human chromosome 7q31. *Nat. Genet.* 27: 392-398.
2. Brown, V.L., et al. 2002. Lack of mutations within ST7 gene in tumour-derived cell lines and primary epithelial tumours. *Br. J. Cancer* 87: 208-211.
3. Dong, S.M. and Sidransky, D. 2002. Absence of ST7 gene alterations in human cancer. *Clin. Cancer Res.* 8: 2939-2941.
4. Vincent, J.B., et al. 2002. The RAY1/ST7 tumor-suppressor locus on chromosome 7q31 represents a complex multi-transcript system. *Genomics* 80: 283-294.
5. Battle, M.A., et al. 2003. ST7 is a novel low-density lipoprotein receptor-related protein (LRP) with a cytoplasmic tail that interacts with proteins related to signal transduction pathways. *Biochemistry* 42: 7270-7282.
6. Wang, S., et al. 2003. An LOH and mutational investigation of the ST7 gene locus in human esophageal carcinoma. *Oncogene* 22: 467-470.
7. Sivasundaram, K., et al. 2003. Mutational analysis of the ST7 gene in human myeloid tumor cell lines. *Oncol. Rep.* 10: 1737-1739.
8. Lu, C., et al. 2003. Somatic mutation analysis of p53 and ST7 tumor suppressor genes in gastric carcinoma by DHPLC. *World J. Gastroenterol.* 9: 2662-2665.

CHROMOSOMAL LOCATION

Genetic locus: ST7 (human) mapping to 7q31.2.

PRODUCT

ST7 (h3): 293T Lysate represents a lysate of human ST7 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

ST7 (h3): 293T Lysate is suitable as a Western Blotting positive control for human reactive ST7 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.