

ephrin-A3 (h3): 293T Lysate: sc-176339

BACKGROUND

The Eph subfamily represents the largest group of receptor protein kinases identified to date. There is increasing evidence that they are involved in central nervous system function and in development. Ligands for Eph receptors include ephrin-A1 (LERK-1/B61), identified as a ligand for the EphA2 (Eck) receptor; ephrin-A2 (ELF-1), identified as a ligand for the EphA3 and EphA4 (Sek) receptors; ephrin-A3 (LERK-3), identified as a ligand for EphA5 (Ehk1) and EphA3 (Hek) receptors; ephrin-A4 (LERK-4), identified as a ligand for the EphA3 receptor; ephrin-A5 (AL-1), identified as a ligand for EphA5 (REK7); ephrin-B1 (LERK-2), identified as a ligand for the EphB1 (Elk) and EphB2 (Cek5) receptors; ephrin-B2 (LERK-5), identified as a ligand for the EphB1, EphB3 (Cek10) and EphB2 receptors; and ephrin-B3 (LERK-8), identified as a ligand for EphB1.

REFERENCES

1. Bartley, T.D., et al. 1994. B61 is a ligand for the Eck receptor protein-tyrosine kinase. *Nature* 368: 558-560.
2. Beckmann, M.P., et al. 1994. Molecular characterization of a family of ligands for Eph-related tyrosine kinase receptors. *EMBO J.* 13: 3757-3762.
3. Cheng, H.J., et al. 1994. Identification and cloning of ELF-1, a developmentally expressed ligand for the MEK-4 and Sek receptor tyrosine kinases. *Cell* 79: 157-168.
4. Kozlosky, C.J., et al. 1995. Ligands for the receptor tyrosine kinases Hek and Elk: isolation of cDNAs encoding a family of proteins. *Oncogene* 10: 299-306.
5. Bergemann, A.D., et al. 1995. ELF-2, a new member of the Eph ligand family, is segmentally expressed in mouse embryos in the region of the hindbrain and newly forming somites. *Mol. Cell. Biol.* 15: 4921-4929.
6. Winslow, J.W., et al. 1995. Cloning of AL-1, a ligand for an Eph-related tyrosine kinase receptor involved in axon bundle formation. *Neuron* 14: 973-981.
7. Gale, N.W., et al. 1996. Elk-LE, a novel transmembrane ligand for the Eph family of receptor tyrosine kinases, expressed in embryonic floor plate, roof plate and hindbrain segments. *Oncogene* 13: 1343-1352.

CHROMOSOMAL LOCATION

Genetic locus: EFNA3 (human) mapping to 1q22.

PRODUCT

ephrin-A3 (h3): 293T Lysate represents a lysate of human ephrin-A3 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.


APPLICATIONS

ephrin-A3 (h3): 293T Lysate is suitable as a Western Blotting positive control for human reactive ephrin-A3 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

ephrin-A3 (12EXZ): sc-73954 is recommended as a positive control antibody for Western Blot analysis of enhanced human ephrin-A3 expression in ephrin-A3 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

DATA



ephrin-A3 (12EXZ): sc-73954. Western blot analysis of ephrin-A3 expression in non-transfected: sc-117752 (A) and human ephrin-A3 transfected: sc-176339 (B) 293T whole cell lysates.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.