

LXR α (h3): 293T Lysate: sc-177491

BACKGROUND

Retinoids are metabolites of vitamin A (retinol) and are believed to represent important signaling molecules during vertebrate development and tissue differentiation. The cooperation of liver X receptors (LXRs) α and β , and retinoic X receptor (RXR) modulate the expression of several genes involved in lipid metabolism in hepatocyte and macrophages. RXR is the receptor for 9-*cis* retinoic acid and dimerizes with VDR, TR, PPAR and several novel receptors including liver X receptors LXR α (also referred to as RLD-1), LXR β and FXR. FXR and LXR fall into a category of proteins termed "orphan receptors" because of their lack of a defined function, and in the case of LXR, the lack of a defined ligand. Both LXR/ RXR and FXR/RXR heterodimers retain their responsiveness to 9-*cis* retinoic acid. LXR α and LXR β share considerable sequence homology and several functions, respond to the same endogenous and synthetic ligands and play critical roles in maintaining lipid homeostasis. LXR β is ubiquitously expressed and enriched in tissues of neuronal and endocrine origin.

REFERENCES

1. Mangelsdorf, D.J., et al. 1994. The retinoid receptors. In Sporn, M.B., et al, eds. The Retinoids: Biology, Chemistry, and Medicine, 2nd Edition. New York: Raven Press, Ltd., 314-349.
2. Bhat, M.K., et al. 1994. Phosphorylation enhances the target gene sequence-dependent dimerization of thyroid hormone receptor with retinoid X receptor. Proc. Natl. Acad. Sci. USA 91: 7927-7931.
3. Song, C., et al. 1994. Ubiquitous receptor: a receptor that modulates gene activation by retinoic acid and thyroid hormone receptors. Proc. Natl. Acad. Sci. USA 91: 10809-10813.
4. Zechel, C., et al. 1994. The dimerization interfaces formed between the DNA binding domains of RXR, RAR and TR determine the binding specificity and polarity of the full-length receptors to direct repeats. EMBO J. 13: 1425-1433.
5. Mangelsdorf, D.J., et al. 1995. The nuclear receptor superfamily: the second decade. Cell 83: 835-839.
6. Mangelsdorf, D.J., et al. 1995. The RXR heterodimers and orphan receptors. Cell 83: 841-850.
7. Willy, P.J., et al. 1995. LXR, a nuclear receptor that defines a distinct retinoid response pathway. Genes Dev. 9: 1033-1045.
8. Leblanc, B.P., et al. 1995. 9-*cis* retinoic acid signaling: changing partners causes some excitement. Genes Dev. 9: 1811-1816.
9. Seol, W.C., et al. 1995. Isolation of proteins that interact specifically with the retinoid X receptor: two novel orphan receptors. Mol. Endocrinol. 9: 72-85.

CHROMOSOMAL LOCATION

Genetic locus: NR1H3 (human) mapping to 11p11.2.

PRODUCT

LXR α (h3): 293T Lysate represents a lysate of human LXR α transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

LXR α (h3): 293T Lysate is suitable as a Western Blotting positive control for human reactive LXR α antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.