

RBM19 (h): 293T Lysate: sc-177849

BACKGROUND

Proteins containing RNA recognition motifs, including various hnRNP proteins, are implicated in the regulation of alternative splicing and protein components of snRNPs. The RBM (RNA-binding motif) gene family encodes proteins with an RNA binding motif that have been suggested to play a role in the modulation of apoptosis. RBM19 (RNA-binding motif protein 19) is a 960 amino acid member of the RRM MRD1 family that contains 6 RRM (RNA recognition motif) domains. Playing a role in embryo pre-implantation development, RBM19 may be involved in regulating ribosome biogenesis. The RBM19 gene is conserved in chimpanzee, canine, bovine, mouse, rat, chicken, zebrafish, fruit fly, mosquito, *C. elegans*, *S. pombe*, *S. cerevisiae*, *K. lactis*, *E. gossypii*, *M. grisea*, *N. crassa*, *A. thaliana*, rice and *P. falciparum*, and maps to human chromosome 12q24.13.

REFERENCES

1. Varani, G. and Nagai, K. 1998. RNA recognition by RNP proteins during RNA processing. *Annu. Rev. Biophys. Biomol. Struct.* 27: 407-445.
2. Mayer, A.N. and Fishman, M.C. 2003. Nil per os encodes a conserved RNA recognition motif protein required for morphogenesis and cytodifferentiation of digestive organs in zebrafish. *Development* 130: 3917-3928.
3. Colland, F., Jacq, X., Trouplin, V., Mougou, C., Groizeleau, C., Hamburger, A., Meil, A., Wojcik, J., Legrain, P. and Gauthier, J.M. 2004. Functional proteomics mapping of a human signaling pathway. *Genome Res.* 14: 1324-1332.
4. Maris, C., Dominguez, C. and Allain, F.H. 2005. The RNA recognition motif, a plastic RNA-binding platform to regulate post-transcriptional gene expression. *FEBS J.* 272: 2118-2131.
5. Lorenzen, J.A., Bonacci, B.B., Palmer, R.E., Wells, C., Zhang, J., Haber, D.A., Goldstein, A.M. and Mayer, A.N. 2005. Rbm19 is a nucleolar protein expressed in crypt/progenitor cells of the intestinal epithelium. *Gene Expr. Patterns* 6: 45-56.
6. Sutherland, L.C., Rintala-Maki, N.D., White, R.D. and Morin, C.D. 2005. RNA binding motif (RBM) proteins: a novel family of apoptosis modulators? *J. Cell. Biochem.* 94: 5-24.
7. Fukuda, T., Naiki, T., Saito, M. and Irie, K. 2009. hnRNP K interacts with RNA binding motif protein 42 and functions in the maintenance of cellular ATP level during stress conditions. *Genes Cells* 14: 113-128.

CHROMOSOMAL LOCATION

Genetic locus: RBM19 (human) mapping to 12q24.13.

PRODUCT

RBM19 (h): 293T Lysate represents a lysate of human RBM19 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

RBM19 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive RBM19 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.