

CD16 (GRM1): sc-19594

BACKGROUND

CD16, the low affinity Fc γ receptor III for IgG (Fc γ RIII), exists both as a polypeptide-anchored form (Fc γ RIIIA or CD16-A) in human natural killer cells and macrophages and as a glycosylphosphatidylinositol-anchored form (Fc γ RIIIB or CD16-B) in neutrophils. CD16-A requires association of the γ subunit of Fc ϵ RI or the ζ subunit of the TCR-CD3 complex for cell surface expression. CD16-B is polymorphic; the two alleles are termed NA1 and NA2. CD16 is one of only four eukaryotic receptors known to exist natively in both the transmembrane (TM, CD16-A) and glycosylphosphatidylinositol (GPI, CD16-B) isoforms. Patients with paroxysmal nocturnal hemoglobinuria (PNH) have only about 10% of the normal levels of CD16 on their neutrophils, whereas the expression of FcRIII is unaffected. Analysis of FcRIII expression in cells of PNH patients, known to be deficient in PI-linked proteins, suggests FcRIII is not PI-linked in monocytes.

REFERENCES

1. Fleit, H.B., et al. 1982. Human neutrophil Fc γ receptor distribution and structure. Proc. Natl. Acad. Sci. USA 79: 3275-3279.
2. Perussia, B., et al. 1984. The Fc receptor for IgG on human natural killer cells: phenotypic, functional, and comparative studies with monoclonal antibodies. J. Immunol. 133: 180-189.
3. Huizinga, T.W., et al. 1988. The PI-linked receptor FcRIII is released on stimulation of neutrophils. Nature 333: 667-669.
4. Nagarajan, S., et al. 1995. Ligand binding and phagocytosis by CD16 (Fc γ receptor III) isoforms. Phagocytic signaling by associated ζ and γ subunits in Chinese hamster ovary cells. J. Biol. Chem. 270: 25762-25770.

CHROMOSOMAL LOCATION

Genetic locus: FCGR3A/FCGR3B (human) mapping to 1q23.3.

SOURCE

CD16 (GRM1) is a mouse monoclonal antibody raised against human prolymphocytic leukemia.

PRODUCT

Each vial contains 200 μ g IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

CD16 (GRM1) is available conjugated to agarose (sc-19594 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-19594 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-19594 PE), fluorescein (sc-19594 FITC), Alexa Fluor[®] 488 (sc-19594 AF488), Alexa Fluor[®] 546 (sc-19594 AF546), Alexa Fluor[®] 594 (sc-19594 AF594) or Alexa Fluor[®] 647 (sc-19594 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-19594 AF680) or Alexa Fluor[®] 790 (sc-19594 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor[®] is a trademark of Molecular Probes, Inc., Oregon, USA

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

CD16 (GRM1) is recommended for detection of CD16 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and flow cytometry (1 μ g per 1 x 10⁶ cells).

Suitable for use as control antibody for CD16 siRNA (h): sc-42758, CD16 shRNA Plasmid (h): sc-42758-SH and CD16 shRNA (h) Lentiviral Particles: sc-42758-V.


Molecular Weight of CD16: 50-100 kDa.

Positive Controls: JM1 whole cell lysate: sc-364233, NK-92 whole cell lysate: 364788 or human spleen extract: sc-363779.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgG κ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



CD16 (GRM1): sc-19594. Western blot analysis of CD16 expression in JM1 whole cell lysate (A) and human spleen tissue extract (B).



CD16 (GRM1): sc-19594. Immunoperoxidase staining of formalin fixed, paraffin-embedded human spleen tissue showing membrane and cytoplasmic staining of cells in white pulp.

SELECT PRODUCT CITATIONS

1. Golay, J., et al. 2019. Human neutrophils express low levels of Fc γ RIIIA, which plays a role in PMN activation. Blood 133: 1395-1405.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.