

CD45R (HIS24): sc-19615

BACKGROUND

CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45R functions as a phosphotyrosine phosphatase, a vital component for efficient tyrosine phosphorylation induction by the TCR/CD3 complex. The tyrosine phosphatase activity of CD45R is contained within the conserved intracellular domain. Src and Syk family protein tyrosine kinases are utilized by the TCR/CD3 complex to initiate signaling cascades. Several members of these two families, including Lck, Fyn and ZAP-70, have been implicated as physiological substrates of CD45R.

CHROMOSOMAL LOCATION

Genetic locus: Ptprc (mouse) mapping to 1 E4.

SOURCE

CD45R (HIS24) is a mouse monoclonal antibody raised against low density cells from AO/G rat Peyer's patches.

PRODUCT

Each vial contains 200 µg IgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

CD45R (HIS24) is available conjugated to either phycoerythrin (sc-19615 PE) or fluorescein (sc-19615 FITC), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

CD45R (HIS24) is recommended for detection of CD45R of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and flow cytometry (1 µg per 1 x 10⁶ cells).

Suitable for use as control antibody for CD45R siRNA (m): sc-35001, CD45R shRNA Plasmid (m): sc-35001-SH and CD45R shRNA (m) Lentiviral Particles: sc-35001-V.


Molecular Weight of CD45R: 240 kDa.

Positive Controls: rat PBL whole cell lysate or rat lymph node tissue extract.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



CD45R (HIS24): sc-19615. Western blot analysis of CD45R expression in rat PBL whole cell lysate (A) and rat lymph node tissue extract (B).



CD45R (HIS24): sc-19615. Immunofluorescence staining of methanol-fixed 3611-RF cells showing membrane localization (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded rat lymph node tissue showing membrane and cytoplasmic staining of cells in germinal center and cells in non-germinal center (B).

SELECT PRODUCT CITATIONS

- Tateda, K., et al. 2012. The suppression of TRIM21 and the accumulation of IFN-α play crucial roles in the pathogenesis of osteonecrosis of the femoral head. *Lab. Invest.* 92: 1318-1329.
- Balint, B., et al. 2019. Left atrial microvascular endothelial dysfunction, myocardial inflammation and fibrosis after selective insular cortex ischemic stroke. *Int. J. Cardiol.* 292: 148-155.
- Yu, J., et al. 2022. Neuron-derived neuropeptide Y fine-tunes the splenic immune responses. *Neuron* 110: 1327-1339.e6.
- Tian, C., et al. 2024. Blocking group 2 innate lymphoid cell activation and macrophage M2 polarization: potential therapeutic mechanisms in ovalbumin-induced allergic asthma by calycosin. *BMC Pharmacol. Toxicol.* 25: 30.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.