# CD83 (HB15E): sc-19678



The Boures to Overtion

#### **BACKGROUND**

CD83 is a heavily glycosylated membrane protein of the immunoglobulin (Ig) superfamily that is expressed in mature dendritic cells, Langerhans cells and interdigitating reticulum cells within lymphoid tissues. Structurally, CD83 resembles other Ig superfamily members, which have an extracellular V-type Ig-like domain, a single transmembrane domain and a 40 residue cytoplasmic tail. CD83 expression is used as a marker for mature, antigen presenting dendritic cells that are capable of generating tumor-specific T cell immunity, a phenotype with implications as an anticancer vaccine. CD83-IgG<sub>1</sub>(fc) chimera studies indicate that CD83 is a sialic acid-binding, Ig-like lectin (Siglec) adhesion molecule that is involved in cell adhesion/signaling by hosting dendritic cell interactions with monocytes and CD8+ T cells.

# **CHROMOSOMAL LOCATION**

Genetic locus: CD83 (human) mapping to 6p23.

#### **SOURCE**

CD83 (HB15E) is a mouse monoclonal antibody raised against COS cells transfected with human CD83.

#### **PRODUCT**

Each vial contains 200  $\mu g \ lg G_1$  kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

CD83 (HB15E) is available conjugated to either phycoerythrin (sc-19678 PE) or fluorescein (sc-19678 FITC), 200  $\mu$ g/ml, for WB (RGB), IF, IHC(P) and FCM.

#### **APPLICATIONS**

CD83 (HB15E) is recommended for detection of CD83 of human origin by immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and flow cytometry (1  $\mu$ g per 1 x 10<sup>6</sup> cells).

Suitable for use as control antibody for CD83 siRNA (h): sc-42808, CD83 shRNA Plasmid (h): sc-42808-SH and CD83 shRNA (h) Lentiviral Particles: sc-42808-V.

Molecular Weight of CD83 precursor: 32 kDa.

Molecular Weight of glycosylated CD83: 45-60 kDa.

# **RECOMMENDED SUPPORT REAGENTS**

To ensure optimal results, the following support reagents are recommended: Immunofluorescence: use m-lgG $\kappa$  BP-FITC: sc-516140 or m-lgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

# **STORAGE**

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.


# **PROTOCOLS**

See our web site at www.scbt.com for detailed protocols and support products.

#### **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

#### DATA



CD83 (HB15E) PE: sc-19678 PE. FCM analysis of K-562human CD83 stable transfectant cell line (solid orange histogram) and non-transfected K-562 (dotted green histogram) cells. Black line histogram represents the isotype control, normal mouse IgG<sub>1</sub>-PE, sc-2866. Transfected cell line kindly provided by Dr. Thomas Tedder at Duke University Medical Center.

# **SELECT PRODUCT CITATIONS**

- 1. Gulubova, M., et al. 2010. Role of TGF- $\beta$ 1, its receptor TGF $\beta$ RII, and Smad proteins in the progression of colorectal cancer. Int. J. Colorectal Dis. 25: 591-599.
- Bühligen, J., et al. 2010. Lysophosphatidylcholine-mediated functional inactivation of syndecan-4 results in decreased adhesion and motility of dendritic cells. J. Cell. Physiol. 225: 905-914.
- Gulubova, M.V., et al. 2012. Role of dendritic cells in progression and clinical outcome of colon cancer. Int. J. Colorectal Dis. 27: 159-169.
- Cirone, M., et al. 2012. Primary effusion lymphoma cell death induced by bortezomib and AG 490 activates dendritic cells through CD91. PLoS ONE 3: e31732.
- Landsverk, O.J., et al. 2012. Differential regulation of MHC II and invariant chain expression during maturation of monocyte-derived dendritic cells. J. Leukoc. Biol. 91: 729-737.
- Gulubova, M., et al. 2013. Relationship of TGF-β1 and Smad7 expression with decreased dendritic cell infiltration in liver gastrointestinal cancer metastasis. APMIS 121: 967-975.
- Bates, J.M., et al. 2015. Dendritic cell CD83 homotypic interactions regulate inflammation and promote mucosal homeostasis. Mucosal Immunol. 8: 414-428.
- 8. Vestre, K., et al. 2021. Rab7b regulates dendritic cell migration by linking lysosomes to the actomyosin cytoskeleton. J. Cell Sci. 134: jcs259221.



See **CD83 (D-3): sc-55536** for CD83 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.