



# cleaved caspase-8 p10 (h385): sc-22180

## BACKGROUND

Initiator caspases, which include caspase-8, activate effector caspases by cleaving inactive forms of effector caspases. In the activation cascade responsible for apoptosis induced by TNFRSF1A and mediated by TNFRSF6/FAS, caspase-8 is the most upstream protease. Caspase-8 binds to adaptor molecule FADD, forming an aggregate referred to as death-inducing signaling complex (DISC). This activates caspase-8. The activated protein is released from the complex and further activates downstream apoptotic proteases. Caspase-8, which is a heterodimer consisting of two subunits (p18 and p10), is widely expressed, but is detected at highest levels in peripheral blood leukocytes (PBLs), thymus, liver and spleen. Defects in CASP8, the gene encoding for caspase-8, may cause CASP8D (caspase-8 deficiency disorder), which is characterized by splenomegaly and CD95-induced apoptosis of PBLs, may lead to immunodeficiency due to defects in T lymphocyte, NK cell and B lymphocyte activation.

## REFERENCES

1. Nagata, S., et al. 1995. The FAS death factor. *Science* 267: 1449-1456.
2. Cleveland, J.L., et al. 1995. Contenders in FAS-L/TNF death signaling. *Cell* 81: 479-482.
3. Fernandes-Alnemri, T., et al. 1996. *In vitro* activation of CPP32 and Mch3 by Mch4, a novel human apoptotic cysteine protease containing two FADD-like domains. *Proc. Natl. Acad. Sci. USA* 93: 7464-7469.
4. Medema, J.P., et al. 1997. FLICE is activated by association with the CD95 death-inducing signaling complex (DISC). *EMBO J.* 16: 2794-804.
5. Srinivasan, A., et al. 1998. Bcl-x<sub>L</sub> functions downstream of caspase-8 to inhibit FAS- and tumor necrosis factor receptor 1-induced apoptosis of MCF7 breast carcinoma cells. *J. Biol. Chem.* 273: 4523-4529.
6. Rytomaa, M., et al. 1999. Involvement of FADD and caspase-8 signalling in detachment-induced apoptosis. *Curr. Biol.* 9: 1043-1046.
7. Wesselborg, S., et al. 1999. Anticancer drugs induce caspase-8/FLICE activation and apoptosis in the absence of CD95 receptor/ligand interaction. *Blood* 93: 3053-3063.
8. Watt, W., et al. 1999. The atomic-resolution structure of human caspase-8, a key activator of apoptosis. *Structure* 7: 1135-1143.
9. Jones, D.T., et al. 2001. Caspase 8 activation independent of FAS (CD95/APO-1) signaling may mediate killing of B-chronic lymphocytic leukemia cells by cytotoxic drugs or  $\gamma$  radiation. *Blood* 98: 2800-2807.

## CHROMOSOMAL LOCATION

Genetic locus: CASP8 (human) mapping to 2q33-q34; Casp8 (mouse) mapping to 1 B.

## SOURCE

cleaved caspase-8 p10 (h385) is a goat polyclonal antibody raised against a short amino acid sequence containing the neoepitope at Leu 385 of caspase-8 p10 of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-22180 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

cleaved caspase-8 p10 (h385) is recommended for detection of the p10 subunit of caspase-8 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with full length caspase-8.

Suitable for use as control antibody for caspase-8 siRNA (h): sc-29930.

cleaved caspase-8 p10: 10 kDa.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.