

E2F-1 Consensus and Mutant Oligonucleotides

BACKGROUND

Electrophoretic mobility shift assays (EMSAs), also known as gel shift assays, provide a relatively straightforward and sensitive method for studying binding interactions between transcription factors and consensus DNA binding elements. For such studies, DNA probes are provided as double-stranded oligonucleotides designed with 5' OH blunt ends to facilitate labeling to high specific activity with polynucleotide kinase. These are constructed both with specific DNA binding consensus sequences for various transcription factors and as control or "mutant" probes in which one or more nucleotides mapping within the consensus binding site has been substituted.

REFERENCES

1. Dignam, J.D., et al. 1983. Accurate transcription initiation by RNA polymerase II in a soluble extract from isolated mammalian nuclei. *Nucleic Acids Res.* 11: 1475-1489.
2. Murre, C., et al. 1991. B cell- and myocyte-specific E2-box-binding factors contain E12/E47-like subunits. *Mol. Cell. Biol.* 11: 1156-1160.
3. Lees, E., et al. 1992. Cyclin E/Cdk2 and cyclin A/Cdk2 kinases associate with p107 and E2F in a temporally distinct manner. *Genes Dev.* 6: 1874-1885.

GEL SHIFT ASSAYS

For gel shift analysis, prepare nuclear extracts following the method of Dignam, et al (1).

- **NOTE:** Spin oligonucleotide vial before opening. Product may be lodged in vial cap.
- Label oligonucleotide probe (TransCruz® Gel Shift Oligonucleotides) with [γ -³²P]-ATP to 50,000 cpm/ng by using polynucleotide kinase.
- Prepare gel shift reaction buffer as follows: 10 mM Tris (Tris: sc-3715), pH 7.5, 50 mM NaCl (NaCl: sc-29108, 1 mM dithiothreitol (DTT: sc-29089), 1 mM EDTA (EDTA: sc-29092), 5% glycerol (glycerol: sc-29095).
- Prepare 20 μ l reaction mixture containing 3-10 μ g nuclear extract and 1 μ g poly dI-dC in gel shift reaction buffer. Add 0.5 ng labeled oligonucleotide probe and incubate for 20 minutes at room temperature. This constitutes the control sample for detection of DNA-protein complexes (2).
- To detect an antibody supershift or block of the DNA-protein complex, prepare reaction mixture as described above, also adding 1-2 μ l of the appropriate TransCruz® Gel Supershift antibody per 20 μ l of reaction volume. Antibody is normally added subsequent to addition of labeled oligonucleotide probe, but result may be improved by adding antibody prior to probe. Incubate at 4° C for 1 hour to overnight, or at room temperature for 15-45 minutes.
- Resolve DNA-protein complexes by electrophoresis (25-35 ma) through a 4% polyacrylamide gel containing 50 mM Tris, pH 7.5, 0.38 M glycine (glycine: sc-29096) and 2 mM EDTA. Dry the gel and visualize by autoradiography.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

E2F-1 CONSENSUS OLIGONUCLEOTIDE: sc-2507

- binding site for E2F-1 transcription factor (3)

```

5' - ATT TAA GTT TCG CGC CCT TTC TCA A - 3'
3' - TAA ATT CAA AGC GCG GGA AAG AGT T - 5'
  
```

E2F-1 MUTANT OLIGONUCLEOTIDE: sc-2508

- identical to sc-2507 with the exception of a "CG" → "AT" substitution in the E2F-1 binding motif (3)

```

5' - ATT TAA GTT TCG ATC CCT TTC TCA A - 3'
3' - TAA ATT CAA AGC TAG GGA AAG AGT T - 5'
  
```

SELECT PRODUCT CITATIONS

1. Canman, C.E., et al. 1998. Small contribution of G₁ checkpoint control manipulation to modulation of p53-mediated apoptosis. *Oncogene* 16: 957-966.
2. Nishikawa, N.S., et al. 2000. Cloning and characterization of the 5'-upstream sequence governing the cell cycle-dependent transcription of mouse DNA polymerase α 68 kDa subunit gene. *Nucleic Acids Res.* 28: 1525-1534.
3. Croxton, R., et al. 2002. Differences in DNA binding properties between E2F1 and E2F4 specify repression of the Mcl-1 promoter. *Oncogene* 21: 1563-1570.
4. Ma, Y., et al. 2003. Regulation of the cyclin D3 promoter by E2F1. *J. Biol. Chem.* 278: 16770-16776.
5. Kwon, J.A., et al. 2003. Transcriptional repression of the human p53 gene by hepatitis B viral core protein (HBc) in human liver cells. *Biol. Chem.* 384: 203-212.
6. Keenan, S.M., et al. 2004. Expression of cyclin E renders cyclin D-Cdk4 dispensable for inactivation of the retinoblastoma tumor suppressor protein, activation of E2F, and G₁-S phase progression. *J. Biol. Chem.* 279: 5387-5396.
7. Sanz-González, S.M., et al. 2004. Role of E2F and ERK1/2 in STI571-mediated smooth muscle cell growth arrest and cyclin A transcriptional repression. *Biochem. Biophys. Res. Commun.* 317: 972-979.
8. Fedele, M., et al. 2011. Expression of a truncated Hmga1b gene induces gigantism, lipomatosis and B-cell lymphomas in mice. *Eur. J. Cancer* 47: 470-478.
9. Hsu, E.C., et al. 2016. Integrin-linked kinase as a novel molecular switch of the IL-6-NF κ B signaling loop in breast cancer. *Carcinogenesis* 37: 430-442.

STORAGE

Store at -20° C; stable for one year from the date of shipment.

NOTE: Spin oligonucleotide vial before opening. Product may be lodged in vial cap.