

RXR (DR-1) mutant oligonucleotide: sc-2548

BACKGROUND

Electrophoretic mobility shift assays (EMSAs), also known as gel shift assays, provide a relatively straightforward and sensitive method for studying binding interactions between transcription factors and consensus DNA binding elements. For such studies, DNA probes are provided as double-stranded oligonucleotides designed with 5' OH blunt ends to facilitate labeling to high specific activity with polynucleotide kinase. These are constructed both with specific DNA binding consensus sequences for various transcription factors and as control or "mutant" probes in which one or more nucleotides mapping within the consensus binding site has been substituted.

REFERENCES

1. Dignam, J.D., et al. 1983. Accurate transcription initiation by RNA polymerase II in a soluble extract from isolated mammalian nuclei. *Nucleic Acids Res.* 11: 1475-1489.
2. Murre, C., et al. 1991. B cell- and myocyte-specific E2-box-binding factors contain E12/E47-like subunits. *Mol. Cell. Biol.* 11: 1156-1160.
3. Kliewer, S.A., et al. 1992. Retinoid X receptor-COUP-TF interactions modulate retinoic acid signaling. *Proc. Natl. Acad. Sci. USA* 89: 1448-1452.

GEL SHIFT ASSAYS

For gel shift analysis, prepare nuclear extracts following the method of Dignam, et al (1).

- **NOTE:** Spin oligonucleotide vial before opening. Product may be lodged in vial cap.
- Label oligonucleotide probe (TransCruz™ Gel Shift Oligonucleotides) with [γ -³²P]-ATP to 50,000 cpm/ng by using polynucleotide kinase.
- Prepare gel shift reaction buffer as follows: 10 mM Tris (Tris: sc-3715), pH 7.5, 50 mM NaCl (NaCl: sc-29108, 1 mM dithiothreitol (DTT: sc-29089), 1 mM EDTA (EDTA: sc-29092), 5% glycerol (glycerol: sc-29095).
- Prepare 20 μ l reaction mixture containing 3-10 μ g nuclear extract and 1 μ g poly dI-dC in gel shift reaction buffer. Add 0.5 ng labeled oligonucleotide probe and incubate for 20 minutes at room temperature. This constitutes the control sample for detection of DNA-protein complexes (2).
- To detect an antibody supershift or block of the DNA-protein complex, prepare reaction mixture as described above, also adding 1-2 μ l of the appropriate TransCruz™ Gel Supershift antibody per 20 μ l of reaction volume. Antibody is normally added subsequent to addition of labeled oligonucleotide probe, but result may be improved by adding antibody prior to probe. Incubate at 4° C for 1 hour to overnight, or at room temperature for 15-45 minutes.
- Resolve DNA-protein complexes by electrophoresis (25-35 ma) through a 4% polyacrylamide gel containing 50 mM Tris, pH 7.5, 0.38 M glycine (glycine: sc-29096) and 2 mM EDTA. Dry the gel and visualize by autoradiography.

PRODUCT

RXR (DR-1) CONSENSUS OLIGONUCLEOTIDE: sc-2547

- binding site for the retinoic X receptor (3)

```

5' - AGC TTC AGG TCA GAG GTC AGA GAG CT - 3'
3' - TCG AAG TCC AGT CTC CAG TCT CTC GA - 5'
  
```

RXR (DR-1) MUTANT OLIGONUCLEOTIDE: sc-2548

- identical to sc-2547 with the exception of two "GT"→"CA" substitutions in the RXR binding motif (3)

```

5' - AGC TTC AGC ACA GAG CAC AGA GAG CT - 3'
3' - TCG AAG TCG TGT CTC GTG TCT CTC GA - 5'
  
```

SELECT PRODUCT CITATIONS

1. Szabo, P.E., et al. 2004. Parent-of-origin-specific binding of nuclear hormone receptor complexes in the H19-IGF-II imprinting control region. *Mol. Cell. Biol.* 24: 4858-4868.
2. Wright, S.K., et al. 2017. Functional activation of PPAR γ in human upper aerodigestive cancer cell lines. *Mol. Carcinog.* 56: 149-162.
3. Rosas, R., et al. 2020. Retinoids augment thiazolidinedione PPAR γ activation in oral cancer cells. *Anticancer Res.* 40: 3071-3080.

STORAGE

Store at -20° C; stable for one year from the date of shipment.

NOTE: Spin oligonucleotide vial before opening. Product may be lodged in vial cap.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.