

Junctophilin-2 siRNA (r): sc-270010

BACKGROUND

Junctophilins are components of the junctional complexes between plasma membranes and endoplasmic or sarcoplasmic reticulum present in all excitable cells. Junctophilins contain a cytoplasmic domain which binds to the plasma membrane, as well as an ER/SR membrane spanning hydrophobic C-terminal segment. The three subtypes in this family are Junctophilin-1, -2 and -3. Junctophilin-1 is predominantly expressed in skeletal muscle, but is also expressed at low levels in heart. Junctophilin-2 is expressed in heart and skeletal muscle. Mutant mice lacking the *Jph2* gene exhibit embryonic lethality and possess cardiac myocytes that express abnormal calcium transients. Junctophilin-3 is expressed in brain. The *JPH3* alternatively spliced exon 2A has been suggested as a site for CTG repeat expansion leading to a Huntington disease-like autosomal dominant disorder.

REFERENCES

1. Takeshima, H., et al. 2000. Junctophilins: a novel family of junctional membrane complex proteins. *Mol. Cell* 6: 11-22.
2. Margolis, R.L., et al. 2001. A disorder similar to Huntington's disease is associated with a novel CAG repeat expansion. *Ann. Neurol.* 50: 373-380.
3. Walker, R.H., et al. 2003. Huntington's disease-like 2 can present as chorea-acanthocytosis. *Neurology* 61: 1002-1004.
4. Margolis, R.L., et al. 2004. Huntington's disease-like 2 (HDL2) in North America and Japan. *Ann. Neurol.* 56: 670-674.
5. Minamisawa, S., et al. 2004. Junctophilin type 2 is associated with caveolin-3 and is downregulated in the hypertrophic and dilated cardiomyopathies. *Biochem. Biophys. Res. Commun.* 325: 852-856.
6. Margolis, R.L., et al. 2005. Huntington's disease-like 2: review and update. *Acta Neurol. Taiwan* 14: 1-8.
7. Matsushita, Y., et al. 2007. Mutation of Junctophilin type 2 associated with hypertrophic cardiomyopathy. *J. Hum. Genet.* 52: 543-548.
8. Rudnicki, D.D., et al. 2007. Huntington's disease-like 2 is associated with CUG repeat-containing RNA foci. *Ann. Neurol.* 61: 272-282.

CHROMOSOMAL LOCATION

Genetic locus: *Jph2* (rat) mapping to 3q42.

PRODUCT

Junctophilin-2 siRNA (r) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Junctophilin-2 shRNA Plasmid (r): sc-270010-SH and Junctophilin-2 shRNA (r) Lentiviral Particles: sc-270010-V as alternate gene silencing products.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Junctophilin-2 siRNA (r) is recommended for the inhibition of Junctophilin-2 expression in rat cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Junctophilin-2 (H-3): sc-377086 is recommended as a control antibody for monitoring of Junctophilin-2 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Junctophilin-2 gene expression knockdown using RT-PCR Primer: Junctophilin-2 (r)-PR: sc-270010-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.