

# Leiomodin 1 siRNA (r): sc-270127

## BACKGROUND

Members of the Leiomodin protein family are closely related to the tropomodulin family of Actin filament pointed end-capping proteins. Leiomodins are characterized as Actin-binding proteins that acts as strong filament nucleators in muscle cells. Leiomodin 1, also known as 64 kDa autoantigen D1 or SM-Lmod, is a 600 amino acid protein that is highly expressed in a variety of tissues that contain smooth muscle and is expressed at lower levels in thyroid and extraocular muscles. Analyzing sera from patients with Hashimoto thyroiditis with thyroid-associated ophthalmopathy (TAO) revealed that antibodies against Leiomodin 1 had been produced in 8 out of 34 patients, while all 12 normal and nonautoimmune individuals were negative. There are two isoforms of Leiomodin 1, which are produced as a result of alternative splicing events.

## REFERENCES

- Dong, Q., Ludgate, M. and Vassart, G. 1991. Cloning and sequencing of a novel 64-kDa autoantigen recognized by patients with autoimmune thyroid disease. *J. Clin. Endocrinol. Metab.* 72: 1375-1381.
- Zhang, Z.G., Dong, Q., Rodien, P., Alcalde, L., Bernard, N., Boucher, A., Salvi, M., Arthurs, B., Vassart, G.M. and Ludgate, M. 1992. Antibodies in the serum of patients with autoimmune thyroid disorders react with a recombinant 98 amino acid fragment of a full length 64 kDa eye muscle membrane protein which is also expressed in the thyroid. *Autoimmunity* 13: 151-157.
- Wall, J.R., Triller, H., Boucher, A., Bernard, N.F., Salvi, M. and Ludgate, M. 1993. Antibodies reactive with an intracellular epitope of a recombinant 64 kDa thyroid and eye muscle protein in patients with thyroid autoimmunity and ophthalmopathy. *J. Endocrinol. Invest.* 16: 863-868.
- Bernard, N.F., Nygen, T.N., Tyutyunikov, A., Stolarski, C., Scalise, D., Genovese, C., Hayes, M.B., Ludgate, M. and Wall, J.R. 1994. Antibodies against 1D, a recombinant 64-kDa membrane protein, are associated with ophthalmopathy in patients with thyroid autoimmunity. *Clin. Immunol. Immunopathol.* 70: 225-233.
- Conley, C.A. and Fowler, V.M. 1999. Localization of the human 64kD autoantigen D1 to myofibrils in a subset of extraocular muscle fibers. *Curr. Eye Res.* 19: 313-322.
- Conley, C.A. 2001. Leiomodin and tropomodulin in smooth muscle. *Am. J. Physiol., Cell Physiol.* 280: C1645-C1656.
- Conley, C.A., Fritz-Six, K.L., Almenar-Queralt, A. and Fowler, V.M. 2001. Leiomodins: larger members of the tropomodulin (Tmod) gene family. *Genomics* 73: 127-139.
- Online Mendelian Inheritance in Man, OMIM™. 2008. Johns Hopkins University, Baltimore, MD. MIM Number: 602715. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## CHROMOSOMAL LOCATION

Genetic locus: Lmod1 (rat) mapping to 13q13.

## PRODUCT

Leiomodin 1 siRNA (r) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Leiomodin 1 shRNA Plasmid (r): sc-270127-SH and Leiomodin 1 shRNA (r) Lentiviral Particles: sc-270127-V as alternate gene silencing products.

For independent verification of Leiomodin 1 (r) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-270127A, sc-270127B and sc-270127C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

Leiomodin 1 siRNA (r) is recommended for the inhibition of Leiomodin 1 expression in rat cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Leiomodin 1 gene expression knockdown using RT-PCR Primer: Leiomodin 1 (r)-PR: sc-270127-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.