

A cyclase I siRNA (chicken): sc-270190

BACKGROUND

Adenylyl cyclases function to convert ATP to cyclic AMP in response to activation by a variety of hormones, neurotransmitters and other regulatory molecules. Cyclic AMP, in turn, activates several other target molecules to control a broad range of diverse phenomena such as metabolism, gene transcription and memory. Adenylyl cyclases respond to receptor-initiated signals, mediated by the G_s and G_i heterotrimeric G proteins. The binding of an agonist to a G_s -coupled receptor catalyzes the exchange of GDP (bound to G_{α_s}) for GTP, the dissociation of $GTP-G_{\alpha_s}$ from $G_{\beta\gamma}$ and G_{α_s} -mediated activation of adenylyl cyclase. A cyclase I, also known as AC1 or ADCY1, is a 1,119 amino acid multi-pass membrane protein expressed in the brain, retina and adrenal medulla. A cyclase I binds two magnesium ions per subunit and may be involved in regulatory processes in the central nervous system.

REFERENCES

- Gilman, A.G. 1987. G proteins: transducers of receptor-generated signals. *Annu. Rev. Biochem.* 56: 615-649.
- Bourne, H.R., Sanders, D.A. and McCormick, F. 1990. The GTPase superfamily: a conserved switch for diverse cell functions. *Nature* 348: 125-132.
- Tang, W.J. and Gilman, A.G. 1992. Adenylyl cyclases. *Cell* 70: 869-872.
- Taussig, R., Tang, W.J., Hepler, J.R. and Gilman, A.G. 1994. Distinct patterns of bidirectional regulation of mammalian adenylyl cyclases. *J. Biol. Chem.* 269: 6093-6100.
- Liu, C.Y., Jamaledin, A.J., Zhang, H. and Christofi, F.L. 1999. FICRHR/cyclic AMP signaling in myenteric ganglia and calbindin-D28 intrinsic primary afferent neurons involves adenylyl cyclases I, III and IV. *Brain Res.* 826: 253-269.
- Gu, C. and Cooper D.M. 2000. Ca^{2+} , Sr^{2+} , and Ba^{2+} identify distinct regulatory sites on adenylyl cyclase (AC) types VI and VIII and consolidate the apposition of capacitative cation entry channels and Ca^{2+} -sensitive ACs. *J. Biol. Chem.* 275: 6980-6986.
- Parkinson, N.A. and Bolsover, S.R. 2001. A nuclear location for Ca^{2+} -activated adenylyl cyclases I and II in neurons. *Brain Res. Mol. Brain Res.* 91: 43-49.
- Scholick, K., Pierre, S. and Patel, T.B. 2001. Protein associated with Myc (PAM) is a potent inhibitor of adenylyl cyclases. *J. Biol. Chem.* 276: 47583-47589.

CHROMOSOMAL LOCATION

Genetic locus: ADCY1 (chicken) mapping to 2.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

A cyclase I siRNA (chicken) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see A cyclase I shRNA Plasmid (chicken): sc-270190-SH and A cyclase I shRNA (chicken) Lentiviral Particles: sc-270190-V as alternate gene silencing products.

For independent verification of A cyclase I (chicken) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-270190A, sc-270190B and sc-270190C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

A cyclase I siRNA (chicken) is recommended for the inhibition of A cyclase I expression in chicken cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor A cyclase I gene expression knockdown using RT-PCR Primer: A cyclase I (chicken)-PR: sc-270190-PR (20 μ l, 485 bp). Annealing temperature for the primers should be $55-60^{\circ}$ C and the extension temperature should be $68-72^{\circ}$ C.