

# T-cadherin siRNA (r): sc-270430

## BACKGROUND

The cadherins are a family of  $\text{Ca}^{2+}$ -dependent adhesion molecules that function to mediate cell-cell binding critical to the maintenance of tissue structure and morphogenesis. Cadherins each contain a large extracellular domain at the amino terminus, which is characterized by a series of five homologous repeats, the most distal of which is thought to be responsible for binding specificity. The relatively short carboxy terminal, intracellular domain interacts with a variety of cytoplasmic proteins, including catenin  $\beta$ , to regulate cadherin function. T-cadherin (for truncated-cadherin, also designated heart-cadherin or cadherin-13) expression levels have been shown to be reduced in human breast cancers and carcinoma cells lines. Evidence suggests that decreased levels of T-cadherin indicate a progression in breast malignancies.

## REFERENCES

1. Takeichi, M. 1988. The cadherins: cell-cell adhesion molecules controlling animal morphogenesis. *Development* 102: 639-655.
2. Hatta, M., et al. 1991. Genomic organization and chromosomal mapping of the mouse P-cadherin gene. *Nucleic Acids Res.* 19: 4437-4441.
3. Hinck, L., et al. 1994. Dynamics of cadherin/catenin complex formation: novel protein interactions and pathways of complex assembly. *J. Cell Biol.* 125: 1327-1340.
4. Ayalon, O., et al. 1994. Spatial and temporal relationships between cadherins and PECAM-1 in cell-cell junctions of human endothelial cells. *J. Cell Biol.* 126: 247-258.
5. Koch, P.J. and Franke, W.W. 1994. Desmosomal cadherins: another growing multigene family of adhesion molecules. *Curr. Opin. Cell Biol.* 6: 682-687.

## CHROMOSOMAL LOCATION

Genetic locus: Cdh13 (rat) mapping to 19q12.

## PRODUCT

T-cadherin siRNA (r) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu\text{M}$  solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see T-cadherin shRNA Plasmid (r): sc-270430-SH and T-cadherin shRNA (r) Lentiviral Particles: sc-270430-V as alternate gene silencing products.

For independent verification of T-cadherin (r) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-270430A, sc-270430B and sc-270430C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at  $-20^{\circ}\text{C}$  with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at  $-20^{\circ}\text{C}$ , avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu\text{l}$  of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu\text{l}$  of RNase-free water makes a 10  $\mu\text{M}$  solution in a 10  $\mu\text{M}$  Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

T-cadherin siRNA (r) is recommended for the inhibition of T-cadherin expression in rat cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu\text{M}$  in 66  $\mu\text{l}$ . Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

T-cadherin (E-9): sc-166875 is recommended as a control antibody for monitoring of T-cadherin gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor T-cadherin gene expression knockdown using RT-PCR Primer: T-cadherin (r)-PR: sc-270430-PR (20  $\mu\text{l}$ ). Annealing temperature for the primers should be  $55-60^{\circ}\text{C}$  and the extension temperature should be  $68-72^{\circ}\text{C}$ .

## SELECT PRODUCT CITATIONS

1. Li, H., et al. 2016. Hyperglycemia abrogates ischemic postconditioning cardioprotection by impairing AdipoR1/Caveolin-3/STAT3 signaling in diabetic rats. *Diabetes* 65: 942-955.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.