



ErbB-3 siRNA (chicken): sc-270447

BACKGROUND

The EGF receptor family comprises several related receptor tyrosine kinases that are frequently overexpressed in a variety of carcinomas. Members of this receptor family include EGFR (HER1), Neu (ErbB-2, HER2), ErbB-3 (HER3) and ErbB-4 (HER4), which form either homodimers or heterodimers upon ligand binding. Full length ErbB-3 is overexpressed in human mammary tumors. The ErbB-3 gene also produces several alternative variants, including a secreted form which negatively regulates heregulin stimulated ErbB activation. ErbB-3 heterodimerizes with Neu and binds heregulin in order to activate phosphoinositide (PI) 3-kinase. The recruitment and activation of PI 3-kinase occurs via its interaction with phosphorylated YXXM motifs in the carboxy terminus of ErbB-3.

REFERENCES

1. Kraus, M.H., et al. 1989. Isolation and characterization of ErbB-3, a third member of the ErbB/epidermal growth factor receptor family: evidence for overexpression in a subset of human mammary tumors. *Proc. Natl. Acad. Sci. USA* 86: 9193-9197.
2. Plowman, G.D., et al. 1990. Molecular cloning and expression of an additional epidermal growth factor receptor-related gene. *Proc. Natl. Acad. Sci. USA* 87: 4905-4909.
3. Kraus, M.H., et al. 1993. Demonstration of ligand-dependent signaling by the ErbB-3 tyrosine kinase and its constitutive activation in human breast tumor cells. *Proc. Natl. Acad. Sci. USA* 90: 2900-2904.
4. Rajkumar, T., et al. 1994. A monoclonal antibody to the human c-ErbB-3 protein stimulates the anchorage-independent growth of breast cancer cell lines. *Br. J. Cancer* 70: 459-465.
5. Rajkumar, T., et al. 1994. The type I growth factor receptors in human breast cancer. *Breast Cancer Res. Treat.* 29: 3-9.
6. Lee, H., et al. 1998. Isolation and characterization of four alternate c-ErbB-3 transcripts expressed in ovarian carcinoma-derived cell lines and normal human tissues. *Oncogene* 16: 3243-3252.
7. Lee, H., et al. 2001. A naturally occurring secreted human ErbB-3 receptor isoform inhibits heregulin-stimulated activation of ErbB-2, ErbB-3 and ErbB-4. *Cancer Res.* 61: 4467-4473.

CHROMOSOMAL LOCATION

Genetic locus: ERBB3 (chicken) mapping to 33.

PRODUCT

ErbB-3 siRNA (chicken) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ErbB-3 shRNA Plasmid (chicken): sc-270447-SH and ErbB-3 shRNA (chicken) Lentiviral Particles: sc-270447-V as alternate gene silencing products.

For independent verification of ErbB-3 (chicken) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-270447A, sc-270447B and sc-270447C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ErbB-3 siRNA (chicken) is recommended for the inhibition of ErbB-3 expression in chicken cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ErbB-3 gene expression knockdown using RT-PCR Primer: ErbB-3 (chicken)-PR: sc-270447-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.