

RPA 70 kDa subunit siRNA (chicken): sc-270630

BACKGROUND

The single-stranded-DNA-binding proteins (SSBs) are essential for DNA function in prokaryotic and eukaryotic cells, mitochondria, phages and viruses. Replication protein A (RPA), a highly conserved eukaryotic protein, is a heterotrimeric SSB. RPA plays an important role in DNA replication, recombination and repair. The binding of human RPA (hRPA) to DNA involves molecular polarity in which initial hRPA binding occurs on the 5' side of an ssDNA substrate and then extends in the 3' direction to create a stably bound hRPA. RPA is a major damage-recognition protein involved in the early stages of nucleotide excision repair. It can also play a role in telomere maintenance. The RPA 70 kDa subunit binds to ssDNA and mediates interactions with many cellular and viral proteins. The DNA binding domain lies in the middle of RPA 70 kDa subunit and comprises two structurally homologous subdomains oriented in tandem. RPA contains a conserved four cysteine-type zinc-finger motif, which mediates the transition of RPA-ssDNA interaction to a stable RPA-ssDNA complex in a redox-dependent manner.

REFERENCES

1. Erdile, L.F., et al. 1990. The primary structure of the 32 kDa subunit of human replication protein A. *J. Biol. Chem.* 265: 3177-3182.
2. Erdile, L.F., et al. 1991. Characterization of a cDNA encoding the 70-kDa single-stranded DNA-binding subunit of human replication protein A and the role of the protein in DNA replication. *J. Biol. Chem.* 266: 12090-12098.
3. Bochkarev, A., et al. 1997. Structure of the single-stranded-DNA-binding domain of replication protein A bound to DNA. *Nature* 385: 176-181.
4. Kim, J., et al. 2000. Replication protein a 32 kDa subunit (RPA p32) binds the SH2 domain of Stat3 and regulates its transcriptional activity. *Cell Biol. Int.* 24: 467-473.
5. Mer, G., et al. 2000. Structural basis for the recognition of DNA repair proteins UNG2, XPA, and Rad52 by replication factor RPA. *Cell* 103: 449-456.
6. Wang, M., et al. 2000. RPA stabilizes the XPA-damaged DNA complex through protein-protein interaction. *Biochemistry* 39: 6433-6439.
7. Iftode, C. and Borowiec, J.A. 2000. 5'→3' molecular polarity of human replication protein (hRPA) binding to pseudo-origin DNA substrates. *Biochemistry* 39: 11970-119981.

PRODUCT

RPA 70 kDa subunit siRNA (chicken) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see RPA 70 kDa subunit shRNA Plasmid (chicken): sc-270630-SH and RPA 70 kDa subunit shRNA (chicken) Lentiviral Particles: sc-270630-V as alternate gene silencing products.

For independent verification of RPA 70 kDa subunit (chicken) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-270630A, sc-270630B and sc-270630C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

RPA 70 kDa subunit siRNA (chicken) is recommended for the inhibition of expression in cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor gene expression knockdown using RT-PCR Primer: RPA 70 kDa subunit (chicken)-PR: sc-270630-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.