

IGF-IIR siRNA (bovine): sc-270658

BACKGROUND

The mannose 6-phosphate/insulin-like growth factor II receptor, IGF-IIR (also designated M6P/IGF2R), is a ubiquitously expressed integral glycoprotein. By binding glycoproteins through two of its extracytoplasmic domains, IGF-IIR mediates the activation of TGF β 1 (a growth inhibitor), the degradation of IGF-I and the transport of lysosomal enzymes. Subsequently, IGF-IIR can form oligomeric complexes, which increase the affinity of IGF-IIR for lysosomal enzymes. Unlike IGF-IR, IGF-IIR does not potentiate the signaling of IGF-I or IGF-II, which have mitogenic, cell survival and Insulin-like effects. Therefore, IGF-IIR is characterized as a tumor suppressor. Furthermore, the IGF-IIR gene is located on chromosome 6q26, which is commonly mutated or deleted in several human cancers.

REFERENCES

1. Ellis, M.J., Jenkins, S., Hanfelt, J., Redington, M.E., Taylor, M., Leek, R., Siddle, K. and Harris, A. 1998. Insulin-like growth factors in human breast cancer. *Breast Cancer Res. Treat.* 52: 175-184.
2. Braulke, T. 1999. Type-2 IGF receptor: a multi-ligand binding protein. *Horm. Metab. Res.* 31: 242-246.
3. Lorenzo, K., Ton, P., Clark, J.L., Coulibaly, S. and Mach, L. 2000. Invasive properties of murine squamous carcinoma cells: secretion of matrix-degrading cathepsins is attributable to a deficiency in the mannose 6-phosphate/Insulin-like growth factor II receptor. *Cancer Res.* 60: 4070-4076.
4. Gemma, A., Hosoya, Y., Uematsu, K., Seike, M., Kurimoto, F., Yoshimura, A., Shibuya, M. and Kudoh, S. 2000. Mutation analysis of the gene encoding the human mannose 6-phosphate/insulin-like growth factor 2 receptor (M6P/IGF2R) in human cell lines resistant to growth inhibition by transforming growth factor β 1 (TGF- β 1). *Lung Cancer* 30: 91-98.
5. Byrd, J.C. and MacDonald, R.G. 2000. Mechanisms for high affinity mannose 6-phosphate ligand binding to the Insulin-like growth factor II/ mannose 6-phosphate receptor. *J. Biol. Chem.* 275: 18638-18646.
6. Byrd, J.C., Park, J.H., Schaffer, B.S., Garmroudi, F. and MacDonald, R.G. 2000. Dimerization of the Insulin-like growth factor II/mannose 6-phosphate receptor. *J. Biol. Chem.* 275: 18647-18656.

CHROMOSOMAL LOCATION

Genetic locus: IGF2R (bovine) mapping to 9q27-q28.

PRODUCT

IGF-IIR siRNA (bovine) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see IGF-IIR shRNA Plasmid (bovine): sc-270658-SH and IGF-IIR shRNA (bovine) Lentiviral Particles: sc-270658-V as alternate gene silencing products.

For independent verification of IGF-IIR (bovine) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-270658A, sc-270658B and sc-270658C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

IGF-IIR siRNA (bovine) is recommended for the inhibition of IGF-IIR expression in bovine cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

IGF-IIR (2G11): sc-53146 is recommended as a control antibody for monitoring of IGF-IIR gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor IGF-IIR gene expression knockdown using RT-PCR Primer: IGF-IIR (bovine)-PR: sc-270658-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.