

SNX4 (B-4): sc-271147

BACKGROUND

Sorting nexin 1 (SNX1) is a member of a large family of hydrophilic proteins that interact with a variety of receptor types and are involved in intracellular trafficking. SNX1 and the related splice variant, SNX1A, bind the epidermal growth factor (EGF) receptor, facilitate its transport to lysosome, and thereby contribute to the degradation of the receptor. SNX2 and SNX4 share a high degree of amino acid similarity with SNX1, as they all contain a characteristic phox homology (PX) domain. These proteins are all partially associated with cellular membranes, and they, likewise, associate with EGF, PDGF and Insulin receptor tyrosine kinases. These nexins are widely expressed and yet have various tissue distribution patterns. Additionally, the sorting nexins can associate with each other and with a variety of other cellular proteins, suggesting that they exist as part of multisubunit complexes. The related protein, SNX3, comprises a distinct subgroup of nexins that share less sequence similarity outside of the PX domain and have dramatically different binding affinities for the tyrosine kinase receptors.

REFERENCES

1. Trowbridge, I.S., et al. 1993. Signal-dependent membrane protein trafficking in the endocytic pathway. *Annu. Rev. Cell Biol.* 9: 129-161.
2. Opresko, L.K., et al. 1995. Endocytosis and lysosomal targeting of epidermal growth factor receptors are mediated by distinct sequences independent of the tyrosine kinase domain. *J. Biol. Chem.* 270: 4325-4333.
3. Kurten, R.C., et al. 1996. Enhanced degradation of EGF receptors by a sorting nexin, SNX1. *Science* 272: 1008-1010.

CHROMOSOMAL LOCATION

Genetic locus: SNX4 (human) mapping to 3q21.2; Snx4 (mouse) mapping to 16 B3.

SOURCE

SNX4 (B-4) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 361-393 within an internal region of SNX4 of human origin.

PRODUCT

Each vial contains 200 µg IgG_{2a} in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

SNX4 (B-4) is available conjugated to agarose (sc-271147 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-271147 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-271147 PE), fluorescein (sc-271147 FITC), Alexa Fluor® 488 (sc-271147 AF488), Alexa Fluor® 546 (sc-271147 AF546), Alexa Fluor® 594 (sc-271147 AF594) or Alexa Fluor® 647 (sc-271147 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-271147 AF680) or Alexa Fluor® 790 (sc-271147 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Blocking peptide available for competition studies, sc-271147 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

SNX4 (B-4) is recommended for detection of SNX4 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


SNX4 (B-4) is also recommended for detection of SNX4 in additional species, including equine and canine.

Suitable for use as control antibody for SNX4 siRNA (h): sc-41353, SNX4 siRNA (m): sc-41354, SNX4 shRNA Plasmid (h): sc-41353-SH, SNX4 shRNA Plasmid (m): sc-41354-SH, SNX4 shRNA (h) Lentiviral Particles: sc-41353-V and SNX4 shRNA (m) Lentiviral Particles: sc-41354-V.


Molecular Weight of SNX4: 60 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204, NIH/3T3 whole cell lysate: sc-2210 or SNX4 (m2): 293T Lysate: sc-123700.

DATA



SNX4 (B-4): sc-271147. Western blot analysis of SNX4 expression in Jurkat (A), HeLa (B), A-431 (C), Hep G2 (D) and NIH/3T3 (E) whole cell lysates.



SNX4 (B-4): sc-271147. Western blot analysis of SNX4 expression in non-transfected: sc-117752 (A) and mouse SNX4 transfected: sc-123700 (B) 293T whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.