

Cdc2 p34 (D-9): sc-271315

BACKGROUND

In vertebrates, as in yeast, multiple cyclins have been identified, including a total of eight such regulatory proteins in mammals. In contrast to the situation in yeast, the Cdc2 p34 kinase is not the only catalytic subunit identified in vertebrates that can interact with cyclins. While Cdc2 p34 is essential for the G₂ to M transition in vertebrate cells, a second Cdc2-related kinase has also been implicated in cell cycle control. This protein, designated cyclin-dependent kinase 2 (Cdk2) p33, also binds to cyclins and its kinase activity is temporally regulated during the cell cycle. Several additional Cdc2 p34-related cyclin dependent kinases have been identified. These include Cdk3-Cdk8, PCTAIRE-1-3 and KIALRE.

CHROMOSOMAL LOCATION

Genetic locus: CDK1 (human) mapping to 10q21.2; Cdk1 (mouse) mapping to 10 B5.3.

SOURCE

Cdc2 p34 (D-9) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 265-297 at the C-terminus of Cdc2 p34 of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-271315 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

Cdc2 p34 (D-9) is recommended for detection of Cdc2 p34 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Cdc2 p34 (D-9) is also recommended for detection of Cdc2 p34 in additional species, including equine and canine.

Suitable for use as control antibody for Cdc2 p34 siRNA (h): sc-29252, Cdc2 p34 siRNA (m): sc-29253, Cdc2 p34 shRNA Plasmid (h): sc-29252-SH, Cdc2 p34 shRNA Plasmid (m): sc-29253-SH, Cdc2 p34 shRNA (h) Lentiviral Particles: sc-29252-V and Cdc2 p34 shRNA (m) Lentiviral Particles: sc-29253-V.


Molecular Weight of Cdc2 p34: 34 kDa.

Positive Controls: MCF7 whole cell lysate: sc-2206, HL-60 whole cell lysate: sc-2209 or K-562 whole cell lysate: sc-2203.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



Cdc2 p34 (D-9): sc-271315. Western blot analysis of Cdc2 p34 expression in HL-60 (A), PC-3 (B), MCF7 (C), BJAB (D) and K-562 (E) whole cell lysates.



Cdc2 p34 (D-9): sc-271315. Immunoperoxidase staining of formalin fixed, paraffin-embedded human testis tissue showing nuclear staining of cells in seminiferous ducts.

SELECT PRODUCT CITATIONS

- Capik, O., et al. 2023. Hypoxia-induced tumor exosomes promote angiogenesis through miR-1825/TSC2/mTOR axis in oral squamous cell carcinoma. Head Neck 45: 2259-2273.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.



See **Cdc2 p34 (17): sc-54** for Cdc2 p34 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.