

Doublecortin (E-6): sc-271390

BACKGROUND

Lissencephaly (smooth brain) is an abnormality of brain development characterized by incomplete neuronal migration and a smooth cerebral surface, resulting in severe mental retardation. Genetic analysis identified two proteins that are mutated in some cases of lissencephaly, designated lissencephaly-1 protein (LIS1) and Doublecortin. LIS1 shows sequence homology to β -subunits of heterotrimeric G proteins. Doublecortin contains a consensus Abl phosphorylation site, and it has some sequence homology to a predicted kinase protein. Both proteins are highly expressed in developing brain, suggesting that they may be involved in a signal transduction pathway that is crucial to brain development.

CHROMOSOMAL LOCATION

Genetic locus: DCX (human) mapping to Xq23, DCLK1 (human) mapping to 13q13.3; Dcx (mouse) mapping to X F2, Dcl1 (mouse) mapping to 3 C.

SOURCE

Doublecortin (E-6) is a mouse monoclonal antibody raised against amino acids 81-365 mapping at the C-terminus of Doublecortin of human origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Doublecortin (E-6) is available conjugated to agarose (sc-271390 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-271390 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-271390 PE), fluorescein (sc-271390 FITC), Alexa Fluor[®] 488 (sc-271390 AF488), Alexa Fluor[®] 546 (sc-271390 AF546), Alexa Fluor[®] 594 (sc-271390 AF594) or Alexa Fluor[®] 647 (sc-271390 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-271390 AF680) or Alexa Fluor[®] 790 (sc-271390 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor[®] is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

Doublecortin (E-6) is recommended for detection of Doublecortin and DCAMK1 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Molecular Weight of Doublecortin: 40 kDa.

Positive Controls: Doublecortin (h): 293T Lysate: sc-114231, SK-N-SH cell lysate: sc-2410 or mouse embryo extract: sc-364239.


STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.


RESEARCH USE

For research use only, not for use in diagnostic procedures.

DATA



Doublecortin (E-6) HRP: sc-271390 HRP. Direct western blot analysis of Doublecortin expression in non-transfected: sc-117752 (A) and human Doublecortin transfected: sc-114231 (B) 293T whole cell lysates and mouse embryo (C) and mouse postnatal brain (D) tissue extracts.



Doublecortin (E-6): sc-271390. Mouse hippocampus (formalin fixed, paraffin) stained with sc-271390; 1:50 overnight with AlexaFluor-568 (red). Green-active caspase-3, Blue-DAPI. Kindly provided by Dr. Svetlana Zonis, Cedars-Sinai Medical Center.

SELECT PRODUCT CITATIONS

- Sirerol-Piquer, M., et al. 2011. GSK3 β overexpression induces neuronal death and a depletion of the neurogenic niches in the dentate gyrus. *Hippocampus* 21: 910-922.
- Kisliouk, T., et al. 2014. Heat stress attenuates new cell generation in the hypothalamus: a role for miR-138. *Neuroscience* 277: 624-636.
- Telias, M., et al. 2015. Molecular mechanisms regulating impaired neurogenesis of fragile X syndrome human embryonic stem cells. *Stem Cells Dev.* 24: 2353-2365.
- García, C.P., et al. 2016. Human pluripotent stem cells and derived neuroprogenitors display differential degrees of susceptibility to BH3 mimetics ABT-263, WEHI-539 and ABT-199. *PLoS ONE* 11: e0152607.
- Agnihotri, S.K., et al. 2017. Loss of PINK1 leads to metabolic deficits in adult neural stem cells and impedes differentiation of newborn neurons in the mouse hippocampus. *FASEB J.* 31: 2839-2853.
- Zonis, S., et al. 2018. Inflammation-induced Gro1 triggers senescence in neuronal progenitors: effects of estradiol. *J. Neuroinflammation* 15: 260.
- Oclair, T., et al. 2019. Mechanisms underpinning AMP-activated protein kinase-related effects on behavior and hippocampal neurogenesis in an animal model of depression. *Neuropharmacology* 150: 121-133.
- Liu, R.X., et al. 2020. No DCX-positive neurogenesis in the cerebral cortex of the adult primate. *Neural Regen. Res.* 15: 1290-1299.
- Wulaer, B., et al. 2021. Shati/Nat8l deficiency disrupts adult neurogenesis and causes attentional impairment through dopaminergic neuronal dysfunction in the dentate gyrus. *J. Neurochem.* 157: 642-655.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.