

mGluR-2 (A-1): sc-271654

BACKGROUND

The mGluR proteins (metabotropic glutamate receptors) are members of the G protein-coupled receptor family and are functionally and pharmacologically distinct from the GluR proteins (ionotropic glutamate receptors). The eight currently known mGluR proteins are mediated by two G-proteins with opposing regulation of adenylate cyclase pathways. The activities of mGluR1 and mGluR5 are mediated by a G-protein that activates a phosphatidylinositol-calcium second messenger system and generates a calcium-activated chloride current. The remainder of the eight sub-types of mGluR have an activity mediated by a G-protein that inhibits adenylate cyclase activity. mGluR-2, which may interact with GRASP, acts as a receptor for glutamate. It may also be involved in the regulation of neurotransmission suppression and in synaptogenesis or synaptic stabilization.

CHROMOSOMAL LOCATION

Genetic locus: GRM2 (human) mapping to 3p21.2; Grm2 (mouse) mapping to 9 F1.

SOURCE

mGluR-2 (A-1) is a mouse monoclonal antibody raised against amino acids 407-536 mapping within an internal region of mGluR-2 of human origin.

PRODUCT

Each vial contains 200 µg IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

mGluR-2 (A-1) is available conjugated to agarose (sc-271654 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-271654 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-271654 PE), fluorescein (sc-271654 FITC), Alexa Fluor® 488 (sc-271654 AF488), Alexa Fluor® 546 (sc-271654 AF546), Alexa Fluor® 594 (sc-271654 AF594) or Alexa Fluor® 647 (sc-271654 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-271654 AF680) or Alexa Fluor® 790 (sc-271654 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

mGluR-2 (A-1) is recommended for detection of mGluR-2 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for mGluR-2 siRNA (h): sc-61028, mGluR-2 siRNA (m): sc-61029, mGluR-2 shRNA Plasmid (h): sc-61028-SH, mGluR-2 shRNA Plasmid (m): sc-61029-SH, mGluR-2 shRNA (h) Lentiviral Particles: sc-61028-V and mGluR-2 shRNA (m) Lentiviral Particles: sc-61029-V.


Molecular Weight of mGluR-2: 110/220 kDa.

Positive Controls: LNCaP cell lysate: sc-2231, IMR-32 cell lysate: sc-2409 or NTERA-2 cl.D1 whole cell lysate: sc-364181.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



mGluR-2 (A-1): sc-271654. Western blot analysis of mGluR-2 expression in LNCaP (A), ZR-75-1 (B), SH-SY5Y (C), F9 (D) and Neuro-2A (E) whole cell lysates.



mGluR-2 (A-1) Alexa Fluor® 488: sc-271654 AF488. Direct fluorescent western blot analysis of mGluR-2 expression in IMR-32 (A), NTERA-2 cl.D1 (B) and U266 (C) whole cell lysates. Blocked with UltraCruz® Blocking Reagent: sc-516214.

SELECT PRODUCT CITATIONS

- Wang, J., et al. 2018. Metabotropic glutamate receptor subtype 2 is a cellular receptor for rabies virus. *PLoS Pathog.* 14: e1007189.
- Wang, J., et al. 2021. SARS-CoV-2 uses metabotropic glutamate receptor subtype 2 as an internalization factor to infect cells. *Cell Discov.* 7: 119.
- Ni, Z., et al. 2024. Influenza virus uses mGluR2 as an endocytic receptor to enter cells. *Nat. Microbiol.* 9: 1764-1777.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.