

# DDX11 (D-2): sc-271711

## BACKGROUND

DEAD-box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp, are putative RNA helicases implicated in several cellular processes involving modifications of RNA secondary structure and ribosome/spliceosome assembly. Based on their distribution patterns, some members of this family may be involved in embryogenesis, spermatogenesis, and cellular growth and division. DDX11 (DEAD/H box protein 11), also known as CHLR1 or KRG2, is a member of the DEAD-box protein family and possesses both ATPase and DNA helicase activity. A homolog of the *S. cerevisiae* CHL1 protein, DDX11 is localized to the nucleus and is highly expressed in the testis, thymus, ovary, spleen and pancreas. DDX11 can bind to both single- and double-stranded DNA and is essential for proper chromosome segregation and embryonic development. Five isoforms of DDX11 exist due to alternative splicing events.

## CHROMOSOMAL LOCATION

Genetic locus: DDX11 (human) mapping to 12p11.21.

## SOURCE

DDX11 (D-2) is a mouse monoclonal antibody raised against amino acids 405-704 mapping within an internal region of DDX11 of human origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>2b</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

DDX11 (D-2) is available conjugated to agarose (sc-271711 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-271711 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-271711 PE), fluorescein (sc-271711 FITC), Alexa Fluor<sup>®</sup> 488 (sc-271711 AF488), Alexa Fluor<sup>®</sup> 546 (sc-271711 AF546), Alexa Fluor<sup>®</sup> 594 (sc-271711 AF594) or Alexa Fluor<sup>®</sup> 647 (sc-271711 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor<sup>®</sup> 680 (sc-271711 AF680) or Alexa Fluor<sup>®</sup> 790 (sc-271711 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

DDX11 (D-2) is recommended for detection of DDX11 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for DDX11 siRNA (h): sc-77104, DDX11 shRNA Plasmid (h): sc-77104-SH and DDX11 shRNA (h) Lentiviral Particles: sc-77104-V.


Molecular Weight of DDX11: 112 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, K-562 whole cell lysate: sc-2203 or Jurkat whole cell lysate: sc-2204.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BPHRP: sc-516102 or m-IgGκ BPHRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, UltraCruz<sup>®</sup> Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BPFITC: sc-516140 or m-IgGκ BPE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz<sup>®</sup> Mounting Medium: sc-24941 or UltraCruz<sup>®</sup> Hard-set Mounting Medium: sc-359850.

## DATA



DDX11 (D-2): sc-271711. Western blot analysis of DDX11 expression in HeLa (A), K-562 (B), SUP-T1 (C) and Jurkat (D) whole cell lysates.



DDX11 (D-2): sc-271711. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic localization. Detection reagent used: goat anti-mouse IgG<sub>2b</sub>-TR: sc-2981 (A). Immunofluorescence staining of methanol-fixed HeLa cells showing nucleolar and nuclear localization (B).

## SELECT PRODUCT CITATIONS

- Cortone, G., et al. 2018. Interaction of the Warsaw breakage syndrome DNA helicase DDX11 with the replication fork-protection factor timeless promotes sister chromatid cohesion. *PLoS Genet.* 14: e1007622.
- Bottega, R., et al. 2019. Two further patients with Warsaw breakage syndrome. Is a mild phenotype possible? *Mol. Genet. Genomic Med.* 7: e639.
- Simon, A.K., et al. 2020. The iron-sulfur helicase DDX11 promotes the generation of single-stranded DNA for CHK1 activation. *Life Sci. Alliance* 3: e201900547.
- Lerner, L.K., et al. 2020. Timeless couples G-quadruplex detection with processing by DDX11 helicase during DNA replication. *EMBO J.* 39: e104185.
- Jegadesan, N.K. and Branzei, D. 2021. DDX11 loss causes replication stress and pharmacologically exploitable DNA repair defects. *Proc. Natl. Acad. Sci. USA* 118: e2024258118.
- Bottega, R., et al. 2021. Genomic integrity and mitochondrial metabolism defects in Warsaw syndrome cells: a comparison with Fanconi anemia. *J. Cell. Physiol.* 236: 5664-5675.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

Alexa Fluor<sup>®</sup> is a trademark of Molecular Probes, Inc., Oregon, USA