

TIMP-2 (H-4): sc-271932

BACKGROUND

TIMP-1, TIMP-2, TIMP-3 and TIMP-4 (for tissue inhibitor of metalloproteinases-1, -2, -3 and -4) complex with metalloproteinases such as collagenases, gelatinases and stromelysins, resulting in irreversible inactivation of the metalloproteinase. TIMP-1 was found to be identical to EPA (erythroid-potentiating activity). Parathyroid hormone has been shown to be a regulator of TIMP-2 in osteoblastic cells. TIMP-3 may be involved in regulating trophoblastic invasion of the uterus as well as in regulating remodeling of the extracellular matrix during the folding of epithelia, and in the formation, branching and expansion of epithelial tubes. TIMP-4 is most highly expressed in heart and low levels of TIMP-4 are expressed in liver, brain, lung, thymus and spleen.

REFERENCES

1. Docherty, A.J., et al. 1985. Sequence of human tissue inhibitor of metalloproteinases and its identity to erythroid-potentiating activity. *Nature* 318: 66-69.
2. Carmichael, D.F., et al. 1986. Primary structure and cDNA cloning of human fibroblast collagenase inhibitor. *Proc. Natl. Acad. Sci. USA* 83: 2407-2411.
3. Cook, T.F., et al. 1994. Cloning and regulation of rat tissue inhibitor of metalloproteinase-2 in osteoblastic cells. *Arch. Biochem. Biophys.* 311: 313-320.
4. Silbiger, S.M., et al. 1994. Cloning of cDNAs encoding human TIMP-3, a novel member of the tissue inhibitor of metalloproteinase family. *Gene* 141: 293-297.
5. Apte, S.S., et al. 1994. Gene encoding a novel murine tissue inhibitor of metalloproteinases (TIMP), TIMP-3, is expressed in developing mouse epithelia, cartilage and muscle, and is located on mouse chromosome 10. *Dev. Dyn.* 200: 177-197.

CHROMOSOMAL LOCATION

Genetic locus: TIMP2 (human) mapping to 17q25.3; Timp2 (mouse) mapping to 11 E2.

SOURCE

TIMP-2 (H-4) is a mouse monoclonal antibody raised against amino acids 81-220 of TIMP-2 of human origin.

PRODUCT

Each vial contains 200 µg IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

TIMP-2 (H-4) is recommended for detection of TIMP-2 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for TIMP-2 siRNA (h): sc-29506, TIMP-2 siRNA (m): sc-37275, TIMP-2 shRNA Plasmid (h): sc-29506-SH, TIMP-2 shRNA Plasmid (m): sc-37275-SH, TIMP-2 shRNA (h) Lentiviral Particles: sc-29506-V and TIMP-2 shRNA (m) Lentiviral Particles: sc-37275-V.


Molecular Weight of TIMP-2: 21 kDa.

Positive Controls: human salivary gland extract: sc-363762, human lung extract: sc-363767 or HeLa whole cell lysate: sc-2200.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



TIMP-2 (H-4): sc-271932. Western blot analysis of human recombinant TIMP-2.



TIMP-2 (H-4): sc-271932. Immunoperoxidase staining of formalin fixed, paraffin-embedded human adrenal gland tissue showing cytoplasmic staining of glandular cells.

SELECT PRODUCT CITATIONS

1. Chaouad, B., et al. 2019. Hyperhomocysteinemia and myocardial remodeling in the sand rat, *Psammomys obesus*. *Acta Histochem.* 121: 823-832.



See **TIMP-2 (3A4): sc-21735** for TIMP-2 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.