

CCL28 (M-19): sc-27341

BACKGROUND

CCL28 functions in chemotactic activity for resting CD4, CD8 T-cells and eosinophils. CCL28, a secreted protein, binds to CCR3 and CCR10 (known previously as orphan G-protein-coupled receptor GPR2) and induces calcium mobilization in a dose-dependent manner. CCL28 is preferentially expressed by epithelial cells of diverse tissues including normal and pathological colon, salivary gland, mammary gland, trachea and rectum. CCL28 belongs to the subfamily of small cytokine CC genes that encode proteins having two adjacent cysteines. Several alternative splice variants may exist.

REFERENCES

1. Wang, W., et al. 2000. Identification of a novel chemokine (CCL28), which binds CCR10 (GPR2). *J. Biol. Chem.* 275: 22313-22323.
2. Lazarus, N.H., et al. 2003. A common mucosal chemokine (mucosae-associated epithelial chemokine/CCL28) selectively attracts IgA plasmablasts. *J. Immunol.* 170: 3799-3805.
3. Hieshima, K., et al. 2003. CCL28 has dual roles in mucosal immunity as a chemokine with broad-spectrum antimicrobial activity. *J. Immunol.* 170: 1452-1461.
4. Wilson, E., et al. 2004. CCL28 controls immunoglobulin (IgA) plasma cell accumulation in the lactating mammary gland and IgA antibody transfer to the neonate. *J. Exp. Med.* 200: 805-809.
5. Hanamoto, H., et al. 2004. Expression of CCL28 by Reed-Sternberg cells defines a major subtype of classical Hodgkin's disease with frequent infiltration of eosinophils and/or plasma cells. *Am. J. Pathol.* 164: 997-1006.
6. SWISS-PROT/TrEMBL ("Q9NRJ3"). World Wide Web URL: <http://www.expasy.ch/sprot/sprot-top.html>.

CHROMOSOMAL LOCATION

Genetic locus: Ccl28 (mouse) mapping to 13.

SOURCE

CCL28 (M-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of CCL28 of mouse origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-27341 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

CCL28 (M-19) is recommended for detection of CCL28 of mouse origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for CCL28 siRNA (m): sc-72145, CCL28 shRNA Plasmid (m): sc-72145-SH and CCL28 shRNA (m) Lentiviral Particles: sc-72145-V.


Molecular Weight of CCL28: 14 kDa.

Positive Controls: CCL28 (m): 293T Lysate: sc-119084.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



CCL28 (M-19): sc-27341. Western blot analysis of CCL28 expression in non-transfected: sc-117752 (A) and mouse CCL28 transfected: sc-119084 (B) 293T whole cell lysates.



CCL28 (M-19): sc-27341. Western blot analysis of mouse recombinant CCL28.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **CCL28 (G-2): sc-376654**, our highly recommended monoclonal alternative to CCL28 (M-19).