

CXCL16 (C-20): sc-27344

BACKGROUND

CXCL16, also known as SR-PSOX (scavenger receptor for phosphatidylserine and oxidized lipoprotein), binds to CXCR6/Bonzo/STRL33, induces a strong chemotactic response and induces calcium mobilization. CXCL16, an α (CXC) chemokine, also has characteristics of C-C chemokines and a structure similar to fractalkine (neurotactin) in having a transmembrane region and a chemokine domain suspended by a mucin-like stalk. Northern blot analysis of mouse and human tissues detect a 2.2-kb CXCL16 transcript in spleen, lymph nodes, Peyer patches, lung, kidney, small intestine and thymus, with weak expression in heart and liver and no expression in brain and bone marrow. Flow cytometry and Western blot analysis demonstrate expression of a glycosylated cell-surface protein and a cell supernatant soluble protein.

REFERENCES

1. Matloubian, M., et al. 2000. A transmembrane C-X-C chemokine is a ligand for HIV-co-receptor Bonzo. *Nat. Immunol.* 1: 298-304.
2. Wilbanks, A., et al. 2001. Expression cloning of the STRL33/Bonzo/TYMSTR ligand reveals elements of C-C, C-X-C, and CX3C chemokines. *J. Immunol.* 166: 5145-5154.
3. Kume, N. 2002. New oxidized LDL receptors and their functions in atherogenesis. *Nippon Ronen Igakkai Zasshi* 39: 264-267.
4. Nakayama, T., et al. 2003. Cutting edge: profile of chemokine receptor expression on human plasma cells accounts for their efficient recruitment to target tissues. *J. Immunol.* 170: 1136-1140.
5. Shashkin, P., et al. 2003. Expression of CXCL16 in human T cells. *Arterioscler. Thromb. Vasc. Biol.* 23: 148-149.
6. Yamauchi, R., et al. 2004. Upregulation of SR-PSOX/CXCL16 and recruitment of CD8⁺ T cells in cardiac valves during inflammatory valvular heart disease. *Arterioscler. Thromb. Vasc. Biol.* 24: 282-287.

CHROMOSOMAL LOCATION

Genetic locus: CXCL16 (human) mapping to 17p13.2.

SOURCE

CXCL16 (C-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of CXCL16 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-27344 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

CXCL16 (C-20) is recommended for detection of CXCL16 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for CXCL16 siRNA (h): sc-105252, CXCL16 shRNA Plasmid (h): sc-105252-SH and CXCL16 shRNA (h) Lentiviral Particles: sc-105252-V.


Molecular Weight of CXCL16: 28 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200 or CXCL16 (h): 293T Lysate: sc-115964.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



CXCL16 (C-20): sc-27344. Western blot analysis of CXCL16 expression in non-transfected: sc-117752 (A) and human CXCL16 transfected: sc-115964 (B) 293T whole cell lysates.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **CXCL16 (C-5): sc-514363** or **CXCL16 (F-7): sc-398792**, our highly recommended monoclonal alternatives to CXCL16 (C-20).