

normal mouse IgG-B: sc-2762

BACKGROUND

Santa Cruz Biotechnology offers a wide variety of control immunoglobulin and control sera for a large selection of species, including mouse, rabbit, goat, chicken, rat, hamster, canine, guinea pig and sheep. Control immunoglobulin and immunoglobulin conjugates are useful negative controls. Normal sera is offered to be used as blocking reagents. Santa Cruz Biotechnology offers affinity purified normal immunoglobulins and immunoglobulin conjugates for use as negative controls in applications including flow cytometry, immunohistochemistry, immunofluorescence, Western Blotting and immunoprecipitation. Agarose (AC) conjugated IgGs are provided for immunoprecipitation; horseradish peroxidase (HRP) conjugates are provided for Western Blotting and immunohistochemistry; and Biotin (B) conjugates are provided for immunohistochemistry. A broad range of fluorescent conjugated controls are also available for use in flow cytometry and immunofluorescence applications. Most control immunoglobulins are available as unconjugated controls or as FITC (fluorescein isothiocyanate), PE (phycoerythrin), PE-Cy5 (phycoerythrin-Cy5), PE-Cy7 (phycoerythrin-Cy7), APC (allophycocyanin) and APC-Cy7 (allophycocyanin-Cy7) conjugates. Additional conjugates include Alexa Fluor[®] 488, Alexa Fluor[®] 647, Alexa Fluor[®] 405, PerCP (peridinin chlorophyll protein complex) and PerCP-Cy5.5 (peridinin chlorophyll protein complex-Cy 5.5). Isotype specific control immunoglobulins include classes such as mouse IgG₁, IgG_{2a}, IgG_{2b}, IgG₃, IgM and IgA, rat IgG₁, IgG_{2a}, IgG_{2b} and IgM, Armenian hamster IgG, and both goat and rabbit IgG.

SOURCE

normal mouse IgG-B is an affinity purified, B (biotin) conjugated isotype control immunoglobulin from mouse.

PRODUCT

Each vial contains 200 µg mouse IgG in 0.5 ml PBS containing 1% protein stabilizer and 0.02% sodium azide.

APPLICATIONS

normal mouse IgG-B is recommended for use as an isotype control immunoglobulin in place of a target specific primary antibody of the same isotype (mouse IgG) by Western Blotting, immunoprecipitation and immunohistochemical staining. To be used at an assay dependent dilution.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

Alexa Fluor[®] is a trademark of Molecular Probes, Inc., Oregon, USA

RECOMMENDED SUPPORT PRODUCTS

- CrystalCruz[™] Cover Glasses, 22 x 50 mm: sc-24975
- PBS, powder: sc-24947
- Formaldehyde: sc-203049
- Hydrogen Peroxide: sc-203336
- Organo/Limonene Mount: sc-45087
- UltraCruz[®] Mounting Medium: sc-24941
- ImmunoHistoMount: sc-45086
- Immuno In Situ Mount: sc-45088
- Paraffin: sc-286633
- Xylenes: sc-237422
- Hematoxylin: sc-24973
- 10X Liquid PBS: sc-24946
- RIPA Lysis Buffer System: sc-24948
- Phenylmethylsulfonyl Fluoride (PMSF): sc-3597
- Electrophoresis Sample Buffer, 2X: sc-24945
- Phosphatase Inhibitor Cocktail A/B: sc-45044/5

SELECT PRODUCT CITATIONS

- Ferraris, J.D., et al. 2002. cAMP-independent role of PKA in tonicity-induced transactivation of tonicity-responsive enhancer/ osmotic response element-binding protein. *Proc. Natl. Acad. Sci. USA* 99: 16800-16805.
- Xu, W.M., et al. 2007. Cystic fibrosis transmembrane conductance regulator is vital to sperm fertilizing capacity and male fertility. *Proc. Natl. Acad. Sci. USA* 104: 9816-9821.
- Silva, M.A., et al. 2008. Intestinal epithelial barrier dysfunction and dendritic cell redistribution during early stages of inflammation in the rat: role for TLR-2 and -4 blockage. *Inflamm. Bowel Dis.* 14: 632-644.
- Tsui, S., et al. 2011. Divergent Sp1 protein levels may underlie differential expression of UDP-glucose dehydrogenase by fibroblasts: role in susceptibility to orbital Graves disease. *J. Biol. Chem.* 286: 24487-24499.
- Guan, W., et al. 2012. S-Nitrosylation of mitogen activated protein kinase phosphatase-1 suppresses radiation-induced apoptosis. *Cancer Lett.* 314: 137-146.
- Chander, H., et al. 2014. Toca-1 is suppressed by p53 to limit breast cancer cell invasion and tumor metastasis. *Breast Cancer Res.* 16: 3413.
- Chen, J., et al. 2016. Soluble egg antigens of *Schistosoma japonicum* induce senescence in activated hepatic stellate cells by activation of the STAT3/p53/p21 pathway. *Sci. Rep.* 6: 30957.

RESEARCH USE

For research use only, not for use in diagnostic procedures.