

EWS (G-5): sc-28327



The Power to Question

BACKGROUND

EWS is a nuclear RNA-binding protein. As a result of chromosome translocation, the EWS gene is fused to a variety of transcription factors, including ATF-1 in human neoplasias. In the Ewing family of tumors, the N-terminal domain of EWS is fused to the DNA-binding domain of various ETS transcription factors, including Fli-1, Erg, ETV1, E1AF and FEV. The EWS/Fli-1 chimeric protein acts as a more potent transcriptional activator than Fli-1 and can promote cell transformation. Two functional regions have been identified in EWS; an amino-terminal region (domain A), that has little transactivation activity but transforms efficiently when fused to Fli-1, and a distal region (domain B), which shows transactivation activity but transforms less efficiently when fused to Fli-1.

REFERENCES

1. Delattre, O., et al. 1992. Gene fusion with an ETS DNA-binding domain caused by chromosome translocation in human tumours. *Nature* 359: 162-165.
2. May, W.A., et al. 1993. The Ewing's sarcoma EWS/Fli-1 fusion gene encodes a more potent transcriptional activator and is a more powerful transforming gene than Fli-1. *Mol. Cell. Biol.* 13: 7393-7398.

CHROMOSOMAL LOCATION

Genetic locus: EWSR1 (human) mapping to 22q12.2; Ewsr1 (mouse) mapping to 11 A1.

SOURCE

EWS (G-5) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 2-19 at the N-terminus of EWS of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

EWS (G-5) is available conjugated to agarose (sc-28327 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-28327 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-28327 PE), fluorescein (sc-28327 FITC), Alexa Fluor® 488 (sc-28327 AF488), Alexa Fluor® 546 (sc-28327 AF546), Alexa Fluor® 594 (sc-28327 AF594) or Alexa Fluor® 647 (sc-28327 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-28327 AF680) or Alexa Fluor® 790 (sc-28327 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Blocking peptide available for competition studies, sc-28327 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

EWS (G-5) is recommended for detection of EWS of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1,000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


EWS (G-5) is also recommended for detection of EWS in additional species, including canine, bovine, porcine and avian.

Suitable for use as control antibody for EWS siRNA (h): sc-35347, EWS siRNA (m): sc-35348, EWS shRNA Plasmid (h): sc-35347-SH, EWS shRNA Plasmid (m): sc-35348-SH, EWS shRNA (h) Lentiviral Particles: sc-35347-V and EWS shRNA (m) Lentiviral Particles: sc-35348-V.

Molecular Weight of EWS: 90 kDa.

Positive Controls: K-562 whole cell lysate: sc-2203, Jurkat whole cell lysate: sc-2204 or F9 cell lysate: sc-2245.

DATA



EWS (G-5) Alexa Fluor® 647: sc-28327 AF647. Direct fluorescent western blot analysis of EWS expression in K-562 (A), F9 (B) and Jurkat (C) whole cell lysates, HeLa nuclear extract (D) and rat testis tissue extract (E). Blocked with UltraCruz® Blocking Reagent: sc-516214.

EWS (G-5): sc-28327. Immunofluorescence staining of methanol-fixed K-562 cells showing nuclear localization (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human colon tissue showing nuclear staining of glandular cells and interstitial cells (B).

SELECT PRODUCT CITATIONS

1. Todd, A.G., et al. 2010. Analysis of SMN-neurite granules: Core Cajal body components are absent from SMN-cytoplasmic complexes. *Biochem. Biophys. Res. Commun.* 397: 479-485.
2. Lindén, M., et al. 2019. FET family fusion oncoproteins target the SWI/SNF chromatin remodeling complex. *EMBO Rep.* 20: e45766.
3. Nicholas, T.R., et al. 2020. —A high-throughput screen identifies inhibitors of the interaction between the oncogenic transcription factor ERG and the cofactor EWS. *PLoS ONE* 15: e0238999.
4. Seong, B.K.A., et al. 2021. TRIM8 modulates the EWS/FLI oncoprotein to promote survival in Ewing sarcoma. *Cancer Cell* 39: 1262-1278.e7.

RESEARCH USE

For research use only, not for use in diagnostic procedures.