SANTA CRUZ BIOTECHNOLOGY, INC.

c-Src siRNA (h): sc-29228



BACKGROUND

The major translational products of the Src gene family are membrane-associated tyrosine protein kinases that lack transmembrane and external amino acid sequences. By virtue of their common structural motifs, the Src family is composed of nine members in vertebrates, including c-Src, c-Yes, Fgr, Yrk, Fyn, Lyn, Hck, Lck and Blk. Src family kinases, which contain an amino-terminal cell membrane anchor followed by SH3 and SH2 domains, transduce signals that are involved in the control of a variety of cellular processes, including proliferation, differentiation, motility and adhesion. Src family members are normally maintained in an inactive state and can be activated transiently during cellular events such as mitosis. Different subcellular locations of Src family kinases may be important for the regulation of specific cellular processes, such as mitogenesis, cytoskeletal organization and membrane trafficking. c-Src (also designated pp60Src, Src p60 and proto-oncogene tyrosine protein kinase Src) is expressed in a broad range of tissue and cell types, although the highest levels of c-Src are detected in neuronal tissues and platelets. c-Src may play a role in events associated with both neuronal differentiation and maintenance of mature neuronal cell functions.

CHROMOSOMAL LOCATION

Genetic locus: SRC (human) mapping to 20q11.23.

PRODUCT

c-Src siRNA (h) is a pool of 4 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see c-Src shRNA Plasmid (h): sc-29228-SH and c-Src shRNA (h) Lentiviral Particles: sc-29228-V as alternate gene silencing products.

For independent verification of c-Src (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-29228A, sc-29228B, sc-29228C and sc-29228D.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNAse-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNAse-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

c-Src siRNA (h) is recommended for the inhibition of c-Src expression in human cells.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

c-Src (H-12): sc-5266 is recommended as a control antibody for monitoring of c-Src gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor c-Src gene expression knockdown using RT-PCR Primer: c-Src (h)-PR: sc-29228-PR (20 μ I, 476 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

DATA





ing of methanol-fixed, control HeLa (A) and c-Src siRNA silenced HeLa (B) cells showing diminished

membrane staining in the siRNA silenced cells. Cells

probed with c-Src (SRC 2): sc-18

c-Src siRNA (h): sc-29228. Western blot analysis of c-Src expression in non-transfected control (Å) and c-Src siRNA transfected (**B**) HeLa cells. Blot probed with c-Src (B-12): sc-8056. GAPDH (FL-335): sc-25778 used as specificity and loading control.

SELECT PRODUCT CITATIONS

- Qiao, F., et al. 2008. Distinct functions of Cdk5(Y15) phosphorylation and Cdk5 activity in stress fiber formation and organization. Exp. Cell Res. 314: 3542-3550.
- 2. Zhu, S., et al. 2019. Activation of IGF1R by DARPP-32 promotes Stat3 signaling in gastric cancer cells. Oncogene 38: 5805-5816.
- Hunter, C.A., et al. 2020. c-Src kinase impairs the expression of mitochondrial OXPHOS complexes in liver cancer. Cell. Signal. 72: 109651.

RESEARCH USE

For research use only, not for use in diagnostic procedures.