

CD34 siRNA (h): sc-29249

BACKGROUND

CD34 is a heavily glycosylated, transmembrane glycoprotein that is expressed on the surface of lymphohematopoietic stem and progenitor cells, small-vessel endothelial cells, embryonic fibroblasts and some cells in fetal and adult nervous tissue. CD34 antigen expression is highest in the most primitive stem cells and is gradually lost as lineage committed progenitors differentiate. The CD34 antigen is also present on capillary endothelial cells and on bone marrow stromal cells. The CD34 cytoplasmic domain has an intracellular domain that contains consensus sites for activated protein kinase C (PKC) phosphorylation as well as serine, threonine and tyrosine phosphorylation consensus sites.

REFERENCES

1. Buck, C.A., et al. 1994. Cell adhesion receptors and early mammalian heart development: an overview. *C.R. Acad. Sci. III* 316: 838-859.
2. DeLisser, H.M., et al. 1994. Platelet endothelial cell adhesion molecule (CD31). *Curr. Top. Microbiol. Immunol.* 184: 37-45.
3. Krause, D.S., et al. 1994. Characterization of murine CD34, a marker for hematopoietic progenitor and stem cells. *Blood* 84: 691-701.
4. Holyoake, T.L. and Alcorn, M.J. 1994. CD34⁺ positive haemopoietic cells: biology and clinical applications. *Blood Rev.* 8: 113-124.
5. Lin, G., et al. 1995. Expression of CD34 in endothelial cells, hematopoietic progenitors and nervous cells in fetal and adult mouse tissues. *Eur. J. Immunol.* 25: 1508-1516.
6. Lasky, L.A. 1995. Selectin-carbohydrate interactions and the initiation of the inflammatory response. *Annu. Rev. Biochem.* 64: 113-139.

CHROMOSOMAL LOCATION

Genetic locus: CD34 (human) mapping to 1q32.2.

PRODUCT

CD34 siRNA (h) is a pool of 4 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see CD34 shRNA Plasmid (h): sc-29249-SH and CD34 shRNA (h) Lentiviral Particles: sc-29249-V as alternate gene silencing products.

For independent verification of CD34 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-29249A, sc-29249B, sc-29249C and sc-29249D.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

CD34 siRNA (h) is recommended for the inhibition of CD34 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.


GENE EXPRESSION MONITORING

CD34 (B-6): sc-74499 is recommended as a control antibody for monitoring of CD34 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor CD34 gene expression knockdown using RT-PCR Primer: CD34 (h)-PR: sc-29249-PR (20 μ l, 405 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

DATA



CD34 siRNA (h): sc-29249. Western blot analysis of CD34 expression in non-transfected control (A) and CD34 siRNA transfected (B) HeLa cells. Blot probed with CD34 (H-140): sc-9095. Lamin A/C (H-110): sc-20681 used as specificity and loading control.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.