

Desmin siRNA (m): sc-29295

BACKGROUND

Cytoskeletal intermediate filaments (IFs) constitute a diverse group of proteins that are expressed in a highly tissue-specific manner. IFs are constructed from two-chain α -helical coiled-coil molecules arranged on an imperfect helical lattice, and have been widely used as markers for distinguishing individual cell types within a tissue and identifying the origins of metastatic tumors. Vimentin is an IF general marker of cells originating in the mesenchyme. Vimentin and Desmin, a related class III IF, are both expressed during skeletal muscle development. Desmin, a 469 amino acid protein found near the Z line in sarcomeres, is expressed more frequently in adult differentiated state tissues. Desmin makes up attachments between the terminal Z-disc and membrane-associated proteins to form a force-transmitting system. Mutations in the gene encoding for Desmin are associated with adult-onset skeletal myopathy, sporadic disease and mild cardiac involvement.

REFERENCES

1. Li, Z.L., et al. 1989. Human Desmin-coding gene: complete nucleotide sequence, characterization and regulation of expression during myogenesis and development. *Gene* 78: 243-254.
2. Tidball, J.G., et al. 1992. Desmin at myotendinous junctions. *Exp. Cell Res.* 199: 206-212.
3. Stewart, M. 1993. Intermediate filament structure and assembly. *Curr. Opin. Cell Biol.* 5: 3-11.
4. Gereben, B., et al. 1995. Species-specificity of glial Vimentin as revealed by immunocytochemical studies with the Vim 3B4 and V9 monoclonal antibodies. *Neurobiology* 3: 151-164.
5. Andreoli, J.M., et al. 1995. Structural and biological consequences of increased Vimentin expression in simple epithelial cell types. *Cell Motil. Cytoskeleton* 32: 10-25.
6. Seshadri, R., et al. 1996. Vimentin expression is not associated with poor prognosis in breast cancer. *Int. J. Cancer* 67: 353-356.

CHROMOSOMAL LOCATION

Genetic locus: Des (mouse) mapping to 1 C4.

PRODUCT

Desmin siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Desmin shRNA Plasmid (m): sc-29295-SH and Desmin shRNA (m) Lentiviral Particles: sc-29295-V as alternate gene silencing products.

For independent verification of Desmin (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-29295A, sc-29295B and sc-29295C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Desmin siRNA (m) is recommended for the inhibition of Desmin expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Desmin (RD301): sc-23879 is recommended as a control antibody for monitoring of Desmin gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Desmin gene expression knockdown using RT-PCR Primer: Desmin (m)-PR: sc-29295-PR (20 μ l, 593 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.