

Pim-3 (4A9): sc-293237

BACKGROUND

The Pim (provirus integration site for Moloney murine leukemia virus) family serine/threonine protein kinases were first identified in studies examining genes targeted for proviral insertion in murine leukemia virus-induced T lymphomas. Increased levels of Pim kinases predispose cells to lymphomagenesis and enhance the activity of mitogenic proteins such as p100, c-Myb and Cdc25A. In addition, Pim kinases are also involved in modulation of synaptic strength in neurons and anti-apoptotic signaling in hematopoietic progenitor cells. Pim-3, a member of the proto-oncogene Pim family that expresses serine/threonine kinase activity, shares significant homology with Pim-1 serine/threonine protein kinases. Pim-3 may function as a mediator of synaptic plasticity in the brain and is presumably involved in the anti-apoptosis process and cell cycle progression as well as the proliferation of human hepatoma cell lines. The Pim-3 protein is widely expressed, however no expression is observed in the colon, thymus or small intestine.

REFERENCES

1. Feldman, J.D., et al. 1998. KID-1, a protein kinase induced by depolarization in brain. *J. Biol. Chem.* 273: 16535-16543.
2. Konietzko, U., et al. 1999. Pim kinase expression is induced by LTP stimulation and required for the consolidation of enduring LTP. *EMBO J.* 18: 3359-3369.
3. Eichmann, A., et al. 2000. Developmental expression of Pim kinases suggests functions also outside of the hematopoietic system. *Oncogene* 19: 1215-1224.
4. Yan, B., et al. 2003. The Pim-2 kinase phosphorylates BAD on Serine 112 and reverses BAD-induced cell death. *J. Biol. Chem.* 278: 45358-45367.
5. Deneen, B., et al. 2003. Pim-3 proto-oncogene kinase is a common transcriptional target of divergent EWS/Ets oncoproteins. *Mol. Cell. Biol.* 23: 3897-3908.
6. Bachmann, M., et al. 2005. The serine/threonine kinase Pim-1. *Int. J. Biochem. Cell Biol.* 37: 726-730.
7. Fujii, C., et al. 2005. Aberrant expression of serine/threonine kinase Pim-3 in hepatocellular carcinoma development and its role in the proliferation of human hepatoma cell lines. *Int. J. Cancer* 114: 209-218.
8. Qian, K.C., et al. 2005. Structural basis of constitutive activity and a unique nucleotide binding mode of human Pim-1 kinase. *J. Biol. Chem.* 280: 6130-6137.
9. Li, Y.Y., et al. 2006. Pim-3, a proto-oncogene with serine/threonine kinase activity, is aberrantly expressed in human pancreatic cancer and phosphorylates BAD to block BAD-mediated apoptosis in human pancreatic cancer cell lines. *Cancer Res.* 66: 6741-6747.

CHROMOSOMAL LOCATION

Genetic locus: PIM3 (human) mapping to 22q13.33.

SOURCE

Pim-3 (4A9) is a mouse monoclonal antibody raised against amino acids 242-326 of Pim-3 of human origin.

PRODUCT

Each vial contains 100 µg IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

Pim-3 (4A9) is recommended for detection of Pim-3 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Suitable for use as control antibody for Pim-3 siRNA (h): sc-61353, Pim-3 shRNA Plasmid (h): sc-61353-SH and Pim-3 shRNA (h) Lentiviral Particles: sc-61353-V.

Molecular Weight of Pim-3: 41 kDa.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

DATA



Pim-3 (4A9): sc-293237. Western blot analysis of human recombinant Pim-3 fusion protein.

SELECT PRODUCT CITATIONS

1. Cho, H., et al. 2020. Anti-survival and pro-apoptotic effects of meridianin C derivatives on MV4-11 human acute myeloid leukemia cells. *Int. J. Oncol.* 56: 368-378.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.