

# HDAC1 siRNA (h): sc-29343

## BACKGROUND

In the intact cell, DNA closely associates with histones and other nuclear proteins to form chromatin. The remodeling of chromatin is believed to be a critical component of transcriptional regulation and a major source of this remodeling is brought about by the acetylation of nucleosomal histones. Acetylation of lysine residues in the amino-terminal tail domain of histone results in an allosteric change in the nucleosomal conformation and an increased accessibility to transcription factors by DNA. Conversely, the deacetylation of histones is associated with transcriptional silencing. Several mammalian proteins have been identified as nuclear histone acetylases, including GCN5, PCAF (for p300/CBP-associated factor), p300/CBP and the TFIID subunit TAF II p250. Mammalian HDAC1 (also designated HD1), HDAC2 (also designated mammalian RPD3) and HDAC3, all of which are related to the yeast transcriptional regulator Rpd3p, have been identified as histone deacetylases.

## CHROMOSOMAL LOCATION

Genetic locus: HDAC1 (human) mapping to 1p35.1.

## PRODUCT

HDAC1 siRNA (h) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see HDAC1 shRNA Plasmid (h): sc-29343-SH and HDAC1 shRNA (h) Lentiviral Particles: sc-29343-V as alternate gene silencing products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

HDAC1 siRNA (h) is recommended for the inhibition of HDAC1 expression in human cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

HDAC1 (10E2): sc-81598 is recommended as a control antibody for monitoring of HDAC1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor HDAC1 gene expression knockdown using RT-PCR Primer: HDAC1 (h)-PR: sc-29343-PR (20  $\mu$ l, 433 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## SELECT PRODUCT CITATIONS

- Chen, C., et al. 2005. Histone acetylation-independent effect of histone deacetylase inhibitors on Akt through the reshuffling of protein phosphatase 1 complexes. *J. Biol. Chem.* 280: 38879-38887.
- Sinha, S., et al. 2010. Coordinated regulation of p53 apoptotic targets Bax and PUMA by SMAR1 through an identical MAR element. *EMBO J.* 29: 830-842.
- Park, S.Y., et al. 2011. Histone deacetylases 1, 6 and 8 are critical for invasion in breast cancer. *Oncol. Rep.* 25: 1677-1681.
- Zhao, H., et al. 2012. The histone acetyltransferase TIP60 interacts with c-Myb and inactivates its transcriptional activity in human leukemia. *J. Biol. Chem.* 287: 925-934.
- Liao, W., et al. 2013. Effect of epigenetic histone modifications on E-cadherin splicing and expression in lung cancer. *Am. J. Cancer Res.* 3: 374-389.
- Liu, C., et al. 2014. Transcriptional repression of the transforming growth factor  $\beta$  (TGF- $\beta$ ) pseudoreceptor BMP and activin membrane-bound inhibitor (BAMBI) by nuclear factor  $\kappa$ B (NF $\kappa$ B) p50 enhances TGF- $\beta$  signaling in hepatic stellate cells. *J. Biol. Chem.* 289: 7082-7091.
- Seo, S.K., et al. 2015. Selective inhibition of histone deacetylase 2 induces p53-dependent survivin downregulation through MDM2 proteasomal degradation. *Oncotarget* 6: 26528-26540.
- Gatla, H.R., et al. 2017. Epigenetic regulation of interleukin-8 expression by class I HDAC and CBP in ovarian cancer cells. *Oncotarget* 8: 70798-70810.
- Li, Y., et al. 2018. Inhibition of miR-34a prevents endothelial cell apoptosis by directly targeting HDAC1 in the setting of atherosclerosis. *Mol. Med. Rep.* 17: 4645-4650.
- Zhou, H., et al. 2019. Aberrant histone deacetylase 1 expression upregulates Vimentin expression via an NF $\kappa$ B-dependent pathway in hepatocellular carcinoma. *Oncol. Lett.* 18: 339-347.
- Bora-Singhal, N., et al. 2020. Novel HDAC11 inhibitors suppress lung adenocarcinoma stem cell self-renewal and overcome drug resistance by suppressing Sox-2. *Sci. Rep.* 10: 4722.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.