

# FGF-8 (2A10): sc-293479

## BACKGROUND

Fibroblast growth factor-1 (FGF-1), also designated acidic FGF, and fibroblast growth factor-2 (FGF-2), also designated basic FGF, are members of a family of growth factors that stimulate proliferation of cells of mesenchymal, epithelial and neuroectodermal origin. Additional members of the FGF family include the oncogenes FGF-3 (Int2) and FGF-4 (hst/Kaposi), FGF-5, FGF-6, FGF-7 (KGF), FGF-8 (AIGF), FGF-9 (GAF) and FGF-10–FGF-23. Members of the FGF family share 30-55% amino acid sequence identity and similar gene structure, and are capable of transforming cultured cells when overexpressed in transfected cells. Cellular receptors for FGFs are members of a second multigene family including four tyrosine kinases, designated Flg (FGFR-1), Bek (FGFR-L), TKF and FGFR-3.

## REFERENCES

- Moore, R., et al. 1986. Sequence, topography and protein coding potential of mouse int-2: a putative oncogene activated by mouse mammary tumor virus. *EMBO J.* 5: 919-924.
- Delli Bovi, P., et al. 1987. An oncogene isolated by transfection of Kaposi's sarcoma DNA encodes a growth factor that is a member of the FGF family. *Cell* 50: 729-737.
- Zhan, X., et al. 1988. The human FGF-5 oncogene encodes a novel protein related to fibroblast growth factors. *Mol. Cell. Biol.* 8: 3487-3495.
- Rifkin, D.B. and Moscatelli, D. 1989. Recent developments in the cell biology of fibroblast growth factor. *J. Cell Biol.* 109: 1-6.
- Marics, I., et al. 1989. Characterization of the HST-related FGF6 gene, a new member of the fibroblast growth factor gene family. *Oncogene* 4: 335-340.
- Tanaka, A., et al. 1992. Cloning and characterization of an androgen-induced growth factor essential for the androgen-dependent growth of mouse mammary carcinoma cells. *Proc. Natl. Acad. Sci. USA* 89: 8928-8932.

## CHROMOSOMAL LOCATION

Genetic locus: FGF8 (human) mapping to 10q24.32; Fgf8 (mouse) mapping to 19 C3.

## SOURCE

FGF-8 (2A10) is a mouse monoclonal antibody raised against amino acids 65-133 representing partial length FGF-8 of human origin.

## PRODUCT

Each vial contains 100 µg IgG<sub>2a</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

FGF-8 (2A10) is recommended for detection of FGF-8 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for FGF-8 siRNA (h): sc-39458, FGF-8 siRNA (m): sc-39459, FGF-8 shRNA Plasmid (h): sc-39458-SH, FGF-8 shRNA Plasmid (m): sc-39459-SH, FGF-8 shRNA (h) Lentiviral Particles: sc-39458-V and FGF-8 shRNA (m) Lentiviral Particles: sc-39459-V.


Molecular Weight of FGF-8: 30 kDa.

Positive Controls: RAW 264.7 whole cell lysate: sc-2211, NIH/3T3 whole cell lysate: sc-2210 or Jurkat whole cell lysate: sc-2204.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

## DATA



FGF-8 (2A10): sc-293479. Western blot analysis of FGF-8 expression in Jurkat (A), NIH/3T3 (B), PC-12 (C) and RAW 264.7 (D) whole cell lysates.



FGF-8 (2A10): sc-293479. Western blot analysis of FGF-8 expression in L8 whole cell lysate.

## SELECT PRODUCT CITATIONS

- Wang, Z., et al. 2021. Overexpression of FGF-8 in the epidermis inhibits hair follicle development. *Exp. Dermatol.* 30: 494-502.
- Lin, C., et al. 2021. FGF-8-mediated signaling regulates tooth developmental pace during odontogenesis. *J. Genet. Genomics.* 49: 40-53.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.