

p21 Waf1/Cip1 siRNA (h): sc-29427

BACKGROUND

It is now well established that cyclins play a positive role in promoting cell cycle transitions via their ability to associate with and activate their cognate cyclin-dependent kinases (Cdks). Cdk2 associates with cyclins A, D and E, and has been implicated in the control of the G₁ to S phase transition in mammals. A novel Cdk-interacting protein, designated p21 Waf1/Cip1, Cip1 or WAF1, has been identified in cyclin A, cyclin D1, cyclin E and Cdk2 immunoprecipitates. p21 Waf1/Cip1 is a potent, tight-binding inhibitor of Cdks and can inhibit the phosphorylation of Rb by cyclin A-Cdk 2, cyclin E-Cdk2, cyclin D1-Cdk4 and cyclin D2-Cdk4 complexes. Expression of p21 Waf1/Cip1 is inducible by wildtype, but not mutant, p53. The mouse homolog of p21 Waf1/Cip1 is designated CAP20.

CHROMOSOMAL LOCATION

Genetic locus: CDKN1A (human) mapping to 6p21.2.

PRODUCT

p21 Waf1/Cip1 siRNA (h) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see p21 Waf1/Cip1 shRNA Plasmid (h): sc-29427-SH and p21 Waf1/Cip1 shRNA (h) Lentiviral Particles: sc-29427-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

p21 Waf1/Cip1 siRNA (h) is recommended for the inhibition of p21 Waf1/Cip1 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

GENE EXPRESSION MONITORING

p21 Waf1/Cip1 (F-5): sc-6246 is recommended as a control antibody for monitoring of p21 Waf1/Cip1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor p21 Waf1/Cip1 gene expression knockdown using RT-PCR Primer: p21 Waf1/Cip1 (h)-PR: sc-29427-PR (20 μ l, 456 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

- Kuo, P.C., et al. 2004. Survivin and p53 modulate quercetin-induced cell growth inhibition and apoptosis in human lung carcinoma cells. *J. Biol. Chem.* 279: 55875-55885.
- Struckhoff, A.P., et al. 2010. Inhibition of p53 sensitizes MCF-7 cells to ceramide treatment. *Int. J. Oncol.* 37: 21-30.
- Braun, F., et al. 2011. Serum-nutrient starvation induces cell death mediated by Bax and Puma that is counteracted by p21 and unmasked by Bcl-x_L inhibition. *PLoS ONE* 6: e23577.
- Kim, D.R., et al. 2012. Combination therapy of conditionally replicating adenovirus and histone deacetylase inhibitors. *Int. J. Mol. Med.* 29: 218-224.
- Parra, E., et al. 2013. Increased expression of p21 Waf1/Cip1 and JNK with costimulation of prostate cancer cell activation by an siRNA Egr-1 inhibitor. *Oncol. Rep.* 30: 911-916.
- Hu, B., et al. 2014. *Ligustrum lucidum* Ait. fruit extract induces apoptosis and cell senescence in human hepatocellular carcinoma cells through upregulation of p21. *Oncol. Rep.* 32: 1037-1042.
- Kim, M.R., et al. 2015. TGF β 1 protects cells from γ -IR by enhancing the activity of the NHEJ repair pathway. *Mol. Cancer Res.* 13: 319-329.
- Kim, D.H., et al. 2016. The insect reptide CopA3 increases colonic epithelial cell proliferation and mucosal barrier function to prevent inflammatory responses in the gut. *J. Biol. Chem.* 291: 3209-3223.
- Das, S., et al. 2017. TRAIL enhances quinacrine-mediated apoptosis in breast cancer cells through induction of autophagy via modulation of p21 and DR5 interactions. *Cell. Oncol.* 40: 593-607.
- Geng, Y., et al. 2018. Trichostatin A promotes GLI1 degradation and p21 expression in multiple myeloma cells. *Cancer Manag. Res.* 10: 2905-2914.
- Waetzig, V., et al. 2019. Retinoic acid-induced survival effects in SH-SY5Y neuroblastoma cells. *J. Cell. Biochem.* 120: 5974-5986.
- Chen, M.C., et al. 2020. RAGE acts as an oncogenic role and promotes the metastasis of human lung cancer. *Cell Death Dis.* 11: 265.

RESEARCH USE

For research use only, not for use in diagnostic procedures.