

Adducin γ siRNA (h): sc-29640

BACKGROUND

Adducins are a family of cytoskeleton proteins encoded by three genes (α , β , γ). Adducin is a protein associated with the inner leaflet of the plasma membrane and is one of the proteins localized at the spectrin-Actin junction of the membrane skeleton. The cortical Actin cytoskeletal network is lost during apoptosis and Adducins are central in the cortical Actin network organization. Adducin α is a cytoskeletal protein involved with sodium-pump activity in the renal tubule and is associated with hypertension. The expression of Adducin α and Adducin γ is ubiquitous in contrast to the restricted expression of Adducin β . Adducin β is expressed at high levels in brain and hematopoietic tissues, such as bone marrow, in humans, and in spleen in mice.

REFERENCES

1. Chapline, C., et al. 1993. Interaction cloning of protein kinase C substrates. *J. Biol. Chem.* 268: 6858-6861.
2. Burns, M.E., et al. 1998. Rabphilin-3A: a multifunctional regulator of synaptic vesicle traffic. *J. Gen. Physiol.* 111: 243-255.
3. Gilligan, D.M., et al. 1999. Targeted disruption of the β Adducin gene (Add2) causes red blood cell spherocytosis in mice. *Proc. Natl. Acad. Sci. USA* 96: 10717-10722.
4. Busjahn, A., et al. 1999. Linkage but lack of association for blood pressure and the α -Adducin locus in normotensive twins. *J. Hypertens.* 17: 1437-1441.

CHROMOSOMAL LOCATION

Genetic locus: ADD3 (human) mapping to 10q25.1.

PRODUCT

Adducin γ siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Adducin γ shRNA Plasmid (h): sc-29640-SH and Adducin γ shRNA (h) Lentiviral Particles: sc-29640-V as alternate gene silencing products.

For independent verification of Adducin γ (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-29640A, sc-29640B and sc-29640C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Adducin γ siRNA (h) is recommended for the inhibition of Adducin γ expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Adducin γ (G-2): sc-365177 is recommended as a control antibody for monitoring of Adducin γ gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Adducin γ gene expression knockdown using RT-PCR Primer: Adducin γ (h)-PR: sc-29640-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

1. Rani, S.B., et al. 2013. MiR-145 functions as a tumor-suppressive RNA by targeting Sox9 and Adducin 3 in human glioma cells. *Neuro Oncol.* 15: 1302-1316.
2. Granato, M., et al. 2015. The activation of KSHV lytic cycle blocks autophagy in PEL cells. *Autophagy* 11: 1978-1986.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.