

Amphiphysin I siRNA (m): sc-29672

BACKGROUND

Amphiphysin is a brain-enriched protein that exhibits N-terminal lipid interaction and functions as a dimer. Amphiphysin contains a membrane bending BAR domain, a middle Clathrin and adaptor binding domain and a C-terminal SH3 domain. In the brain, Amphiphysin I and II form heterodimers that bind to the Clathrin-associated GTPase Dynamin via their SH3 domains. This association is essential for synaptic vesicle recycling in neurons, as it precedes the binding of Dynamin to the Clathrin-coated pits and the subsequent vesicle budding. In other tissues, Amphiphysin may play a key role in other membrane bending and curvature stabilization events. The mammalian Amphiphysins, Amphiphysin I and Amphiphysin II, have similar overall structure. An ubiquitous splice form of Amphiphysin II that does not contain Clathrin or adaptor interactions is highly expressed in muscle tissue and is involved in the formation and stabilization of the T tubule network.

REFERENCES

1. Lichte, B., et al. 1992. Amphiphysin, a novel protein associated with synaptic vesicles. *EMBO J.* 11: 2521-2530.
2. Yamamoto, R., et al. 1995. Primary structure of human amphiphysin, the dominant autoantigen of paraneoplastic stiff-man syndrome, and mapping of its gene (AMPH) to chromosome 7p13-p14. *Hum. Mol. Genet.* 4: 265-268.
3. Sakamuro, D., et al. 1996. BIN1 is a novel Myc-interacting protein with features of a tumour suppressor. *Nat. Genet.* 14: 69-77.
4. Leprince, C., et al. 1997. A new member of the amphiphysin family connecting endocytosis and signal transduction pathways. *J. Biol. Chem.* 272: 15101-15105.
5. Wechsler-Reya, R., et al. 1997. Structural analysis of the human BIN1 gene. Evidence for tissue-specific transcriptional regulation and alternate RNA splicing. *J. Biol. Chem.* 272: 31453-31458.
6. Wigge, P., et al. 1997. Amphiphysin heterodimers: potential role in Clathrin-mediated endocytosis. *Mol. Biol. Cell* 8: 2004-2015.

CHROMOSOMAL LOCATION

Genetic locus: Amph (mouse) mapping to 13 A2.

PRODUCT

Amphiphysin I siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Amphiphysin I shRNA Plasmid (m): sc-29672-SH and Amphiphysin I shRNA (m) Lentiviral Particles: sc-29672-V as alternate gene silencing products.

For independent verification of Amphiphysin I (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-29672A, sc-29672B and sc-29672C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Amphiphysin I siRNA (m) is recommended for the inhibition of Amphiphysin I expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Amphiphysin I (G-4): sc-376402 is recommended as a control antibody for monitoring of Amphiphysin I gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Amphiphysin I gene expression knockdown using RT-PCR Primer: Amphiphysin I (m)-PR: sc-29672-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.