

CAS siRNA (m): sc-29909

BACKGROUND

Cellular apoptosis susceptibility protein (CAS), also called Exportin 2, is a 971 amino acid member of the CSE1 family. CAS mediates Importin α re-export from the nucleus to the cytoplasm after import substrates have been released into the nucleoplasm. In the nucleus, CAS binds cooperatively to Importin α and to the GTPase Ran in its GTP-bound (active) form. This complex binds to nucleoporins as it docks to the nuclear pore complex. Once in the cytoplasm, the complex dissociates and Importin α is released and CAS returns to the nuclear compartment and the process begins anew. CAS can be detected highly in proliferating cells. Three isoforms of CAS have been named due to alternative splicing. Isoform 1 is the full length, 971 amino acid protein. Isoform 2 contains an alternative sequence for amino acids 190-195 and is missing amino acids 196-971. Isoform 3 contains an alternative sequence for amino acids 943-945 and is missing amino acids 946-971.

REFERENCES

1. Brinkmann, U., et al. 1995. Cloning and characterization of a cellular apoptosis susceptibility gene, the human homologue to the yeast chromosome segregation gene CSE1. *Proc. Natl. Acad. Sci. USA* 92: 10427-10431.
2. Columbano, A. 1995. Cell death: current difficulties in discriminating apoptosis from necrosis in the context of pathological processes *in vivo*. *J. Cell. Biochem.* 58: 181-190.

CHROMOSOMAL LOCATION

Genetic locus: Cse1l (mouse) mapping to 2 H3.

PRODUCT

CAS siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see CAS shRNA Plasmid (m): sc-29909-SH and CAS shRNA (m) Lentiviral Particles: sc-29909-V as alternate gene silencing products.

For independent verification of CAS (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-29909A, sc-29909B and sc-29909C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

APPLICATIONS

CAS siRNA (m) is recommended for the inhibition of CAS expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING


CAS (H-2): sc-271537 is recommended as a control antibody for monitoring of CAS gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz MarkerTM Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor CAS gene expression knockdown using RT-PCR Primer: CAS (m)-PR: sc-29909-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

DATA



CAS siRNA (m): sc-29909. Western blot analysis of CAS expression in non-transfected control (A) and CAS siRNA transfected (B) NIH/3T3 cells. Blot probed with CAS (N-19): sc-1709. Actin (I-19): sc-1616 used as specificity and loading control.

RESEARCH USE

For research use only, not for use in diagnostic procedures.