

CD36 siRNA (h): sc-29995

BACKGROUND

CD36 (collagen type I receptor, thrombospondin receptor, FAT, GP4, GP3B, GPIV, PASIV, SCARB3) is a membrane glycoprotein on platelets, monocytes and umbilical vein endothelial cells. CD36 binds to collagen, thrombospondin, anionic phospholipids and oxidized LDL. CD36 plays a key role in both phagocytosis and lipid recycling, for constant production of mature spermatozoa. Mutations in this gene cause platelet glycoprotein deficiency. Three alternatively spliced transcript variants encoding the same protein isoform have been found for this gene. Thrombospondins are widely distributed proteins that influence a variety of adhesive processes and CD36 may have important functions as a cell adhesion molecule.

CHROMOSOMAL LOCATION

Genetic locus: CD36 (human) mapping to 7q21.11.

PRODUCT

CD36 siRNA (h) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see CD36 shRNA Plasmid (h): sc-29995-SH and CD36 shRNA (h) Lentiviral Particles: sc-29995-V as alternate gene silencing products.

For independent verification of CD36 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-29995A and sc-29995B.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

CD36 siRNA (h) is recommended for the inhibition of CD36 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

CD36 (SM ϕ): sc-7309 is recommended as a control antibody for monitoring of CD36 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor CD36 gene expression knockdown using RT-PCR Primer: CD36 (h)-PR: sc-29995-PR (20 μ l, 524 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

- Subramaniam, D., et al. 2010. Cholesterol rich lipid raft microdomains are gateway for acute phase protein, SERPINA1. *Int. J. Biochem. Cell Biol.* 42: 1562-1570.
- Zhou, M.S., et al. 2013. Nicotine potentiates proatherogenic effects of oxLDL by stimulating and upregulating macrophage CD36 signaling. *Am. J. Physiol. Heart Circ. Physiol.* 305: H563-H574.
- Gao, X., et al. 2014. Oxidized high-density lipoprotein impairs the function of human renal proximal tubule epithelial cells through CD36. *Int. J. Mol. Med.* 34: 564-572.
- Roggia, M.F. and Ueta, T. 2015. $\alpha_v\beta_5$ Integrin/FAK/PGC-1 α pathway confers protective effects on retinal pigment epithelium. *PLoS ONE* 10: e0134870.
- Hou, Y., et al. 2015. CD36 is involved in high glucose-induced epithelial to mesenchymal transition in renal tubular epithelial cells. *Biochem. Biophys. Res. Commun.* 468: 281-286.
- Scipione, C.A., et al. 2015. Mechanistic insights into Lp(a)-induced IL-8 expression: a role for oxidized phospholipid modification of apo(a). *J. Lipid Res.* 56: 2273-2285.
- Cheng, J.J., et al. 2016. CD36 is a co-receptor for hepatitis C virus E1 protein attachment. *Sci. Rep.* 6: 21808.
- Choi, Y.J., et al. 2017. Activation of AMPK by berberine induces hepatic lipid accumulation by upregulation of fatty acid translocase CD36 in mice. *Toxicol. Appl. Pharmacol.* 316: 74-82.
- Dai, Y., et al. 2017. Xanthine Oxidase induces foam cell formation through LOX-1 and NLRP3 activation. *Cardiovasc. Drugs Ther.* 31: 19-27.
- Madak-Erdogan, Z., et al. 2019. Free fatty acids rewire cancer metabolism in obesity-associated breast cancer via estrogen receptor and mTOR signaling. *Cancer Res.* 79: 2494-2510.
- Ding, D., et al. 2019. MicroRNA-26a-CD36 signaling pathway: pivotal role in lipid accumulation in hepatocytes induced by PM2.5 liposoluble extracts. *Environ. Pollut.* 248: 269-278.

RESEARCH USE

For research use only, not for use in diagnostic procedures.