

# cystatin B (RJMW 2E7): sc-32802

## BACKGROUND

Cystatin A (also designated STF1, STFA, stefin A or cystatin AS) and cystatin B (also designated PME, CST6, STFB, CPI-B, stefin B and liver thiol proteinase inhibitor) are thiol protease inhibitors that form complexes with papain and the cathepsins B, H, and L. Cystatin A, a cytoplasmic protein, is one of the precursor proteins of the cornified cell envelope in keratinocytes and plays a role in epidermal development and maintenance. Cystatin B protects against intracellular proteases leaking out of lysosomes and is primarily expressed in heart, liver and kidney.

## REFERENCES

- Ritonja, A., et al. 1985. Amino acid sequence of the intracellular cysteine proteinase inhibitor cystatin B from human liver. *Biochem. Biophys. Res. Commun.* 131: 1187-1192.
- Jerala, R., et al. 1988. Cloning a synthetic gene for human stefin B and its expression in *E. coli*. *FEBS Lett.* 239: 41-44.
- Pennacchio, L.A., et al. 1996. Mutations in the gene encoding cystatin B in progressive myoclonus epilepsy (EPM1). *Science* 271: 1731-1734.
- Kos, J., et al. 1998. Cysteine proteinases and their endogenous inhibitors: target proteins for prognosis, diagnosis and therapy in cancer (review). *Oncol. Rep.* 5: 1349-1361.
- Takahashi, H., et al. 1998. Structure and transcriptional regulation of the human cystatin A gene. The 12-O-tetradecanoylphorbol-13-acetate (TPA) responsive element-2 site (-272 to -278) on cystatin A gene is critical for TPA-dependent regulation. *J. Biol. Chem.* 273: 17375-17380.
- Takahashi, H., et al. 2001. Expression of human cystatin A by keratinocytes is positively regulated via the Ras/MEK1/MKK7/JNK signal transduction pathway but negatively regulated via the Ras/Raf-1/MEK1/ERK pathway. *J. Biol. Chem.* 276: 36632-36638.
- Jenko, S., et al. 2004. Different propensity to form amyloid fibrils by two homologous proteins—human stefins A and B: searching for an explanation. *Proteins* 55: 417-425.

## CHROMOSOMAL LOCATION

Genetic locus: CSTB (human) mapping to 21q22.3.

## SOURCE

cystatin B (RJMW 2E7) is a mouse monoclonal antibody raised against human spleen cystatin B.

## PRODUCT

Each vial contains 200 µg IgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

cystatin B (RJMW 2E7) is recommended for detection of cystatin B of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

cystatin B (RJMW 2E7) is also recommended for detection of cystatin B in additional species, including monkey.

Suitable for use as control antibody for cystatin B siRNA (h): sc-44431, cystatin B shRNA Plasmid (h): sc-44431-SH and cystatin B shRNA (h) Lentiviral Particles: sc-44431-V


Molecular Weight of cystatin B: 12 kDa.

Positive Controls: cystatin B (h2): 293 Lysate: sc-112337, U-87 MG cell lysate: sc-2411 or T98G cell lysate: sc-2294.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:  
 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.  
 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).  
 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## DATA



cystatin B (RJMW 2E7): sc-32802. Western blot analysis of cystatin B expression in U-87 MG (A), T98G (B), SK-N-MC (C) and WI-38 (D) whole cell lysates.



cystatin B (RJMW 2E7): sc-32802. Western blot analysis of cystatin B expression in non-transfected 293: sc-110760 (A), human cystatin B transfected 293: sc-112337 (B) and U-87 MG (C) whole cell lysates.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.