

IFN- γ (3F1E3): sc-32813

BACKGROUND

Interferon (IFN)- γ is an antiviral and antiparasitic agent produced by CD4⁺/CD8⁺ lymphocytes and natural killer cells that undergo activation by antigens, mitogens or alloantigens. IFN- γ production modulates T cell growth and differentiation and inhibits the growth of B cells. Synthesis of IFN- γ is inducible by IL-2, FGF and EGF. The active form of IFN- γ is a homodimer with each subunit containing six helices. The dimeric structure of human IFN- γ is stabilized by non-covalent interactions through the interface of the helices. IFN- γ translated precursor is 166 amino acids, including the 23 amino acid secretory sequence. 20 kDa and 25 kDa forms exist due to variable glycosylation; 40-60 kDa forms are observable under non-denaturing conditions as dimers and tetramers.

REFERENCES

- Young, H.A., et al. 1995. Role of IFN- γ in immune cell regulation. *J. Leukoc. Biol.* 58: 373-381.
- Dinarello, C.A., et al. 1998. Overview of interleukin-18: more than an IFN- γ inducing factor. *J. Leukoc. Biol.* 63: 658-664.
- Okamura, H., et al. 1998. Regulation of IFN- γ production by IL-12 and IL-18. *Curr. Opin. Immunol.* 10: 259-264.
- Costa-Pereira, A.P., et al. 2002. The antiviral response to IFN- γ . *J. Virol.* 76: 9060-9068.
- Zika, E., et al. 2003. Histone deacetylase 1/mSin3A disrupts IFN- γ -induced CIITA function and major histocompatibility complex class II enhanceosome formation. *Mol. Cell. Biol.* 23: 3091-3102.
- Schroder, K., et al. 2004. IFN- γ : an overview of signals, mechanisms and functions. *J. Leukoc. Biol.* 75: 163-189.
- Ellis, T.N., et al. 2004. IFN- γ activation of polymorphonuclear neutrophil function. *Immunology* 112: 2-12.
- Sizemore, N., et al. 2004. Inhibitor of κ B kinase is required to activate a subset of interferon- γ -stimulated genes. *Proc. Natl. Acad. Sci. USA* 101: 7994-7998.
- Halfter, U.M., et al. 2005. Interferon- γ -dependent tyrosine phosphorylation of MEKK-4 via Pyk2 is regulated by annexin II and SHP2 in keratinocytes. *Biochem. J.* 388: 17-28.

CHROMOSOMAL LOCATION

Genetic locus: IFNG (human) mapping to 12q15.

SOURCE

IFN- γ (3F1E3) is a mouse monoclonal antibody raised against recombinant human IFN- γ .

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

IFN- γ (3F1E3) is recommended for detection of precursor and mature IFN- γ of human origin by Western Blotting (starting dilution 1:100, dilution range) and immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)].

Suitable for use as control antibody for IFN- γ siRNA (h): sc-39606, IFN- γ shRNA Plasmid (h): sc-39606-SH and IFN- γ shRNA (h) Lentiviral Particles: sc-39606-V.

Molecular Weight of IFN- γ : 20-25 kDa.


Positive Controls: Jurkat whole cell lysate: sc-2204, CCRF-CEM cell lysate: sc-2225 or Daudi cell lysate: sc-2415.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:

1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

DATA



IFN- γ (3F1E3): sc-32813. Western blot analysis of human recombinant IFN- γ .

SELECT PRODUCT CITATIONS

- Zhou, X., et al. 2015. Transient receptor potential channel 1 deficiency impairs host defense and proinflammatory responses to bacterial infection by regulating protein kinase α signaling. *Mol. Cell. Biol.* 35: 2729-2739.
- Eskicioglu, F., et al. 2015. The association of HLA-G and immune markers in recurrent miscarriages. *J. Matern. Fetal Neonatal Med.* 29: 3056-3060.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.



See **IFN- γ (E-10): sc-373727** for IFN- γ antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.