SANTA CRUZ BIOTECHNOLOGY, INC.

GAP-43 (7B10): sc-33705



BACKGROUND

GAP-43 (growth associated protein 43, B-50, PP46, calmodulin-binding protein P-57, neuromodulin, neuron growth-associated protein 43, protein F1) is a crucial component for regenerative response in the nervous system that is present at high levels in neuronal growth cones during development and axonal regeneration. GAP-43 is normally produced by neurons during developmental growth and axonal regeneration, but it is also expressed in specific regions of the normal adult nervous system. The neuron-specific Elav/Hu family member, HuD, interacts with and stabilizes GAP-43 mRNA in developing neurons and leads to increased levels of GAP-43 protein. Heterozygous GAP-43 knockout mice with GAP-43 levels reduced by onehalf display significant memory impairments in cued conditioning or on tests of nociceptive or auditory perception.

CHROMOSOMAL LOCATION

Genetic locus: GAP43 (human) mapping to 3q13.31; Gap43 (mouse) mapping to 16 B4.

SOURCE

GAP-43 (7B10) is a mouse monoclonal antibody raised against full length GAP-43 of rat origin.

PRODUCT

Each vial contains 200 μg IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

GAP-43 (7B10) is available conjugated to agarose (sc-33705 AC), 500 μ g/ 0.25 ml agarose in 1 ml, for IP; to HRP (sc-33705 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-33705 PE), fluorescein (sc-33705 FITC), Alexa Fluor* 488 (sc-33705 AF488), Alexa Fluor* 546 (sc-33705 AF546), Alexa Fluor* 594 (sc-33705 AF594) or Alexa Fluor* 647 (sc-33705 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor* 680 (sc-33705 AF680) or Alexa Fluor* 790 (sc-33705 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

GAP-43 (7B10) is recommended for detection of axonal membrane protein GAP-43 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for GAP-43 siRNA (h): sc-35446, GAP-43 siRNA (m): sc-35447, GAP-43 shRNA Plasmid (h): sc-35446-SH, GAP-43 shRNA Plasmid (m): sc-35447-SH, GAP-43 shRNA (h) Lentiviral Particles: sc-35446-V and GAP-43 shRNA (m) Lentiviral Particles: sc-35447-V.

Molecular Weight of GAP-43: 43 kDa.

Positive Controls: mouse spinal cord extract: sc-395045, rat cerebellum extract: sc-2398 or rat brain extract: sc-2392.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

DATA





GAP-43 (7B10): sc-33705. Western blot analysis of GAP-43 expression in human brain (A), mouse cerebellum (B), mouse spinal cord (C), rat brain (D), rat cerebellum (E) and rat spinal cord (F) tissue extracts Detection reagent used: m-lgG Fc BP-HRP: sc-525409.

GAP-43 (7B10): sc-33705. Immunoperoxidase staining of formalin fixed, paraffin-embedded human cerebral cortex (A) and mouse brain (B) tissue showing neuropil staining.

SELECT PRODUCT CITATIONS

- Teunissen, C.E., et al. 2006. Growth-associated protein 43 in lesions and cerebrospinal fluid in multiple sclerosis. Neuropathol. Appl. Neurobiol. 32: 318-331.
- Lemmens, K., et al. 2016. Matrix metalloproteinases as promising regulators of axonal regrowth in the injured adult zebrafish retinotectal system. J. Comp. Neurol. 524: 1472-1493.
- 3. Zhang, X., et al. 2017. MicroRNA-210 contributes to peripheral nerve regeneration through promoting the proliferation and migration of Schwann cells. Exp. Ther. Med. 14: 2809-2816.
- 4. Shi, D., et al. 2018. Local application of MDL28170-loaded PCL film improves functional recovery by preserving survival of motor neurons after traumatic spinal cord injury. Neurosci. Lett. 694: 161-167.
- Zhang, K., et al. 2019. ALK5 signaling pathway mediates neurogenesis and functional recovery after cerebral ischemia/reperfusion in rats via Gadd45b. Cell Death Dis. 10: 360.
- Afzal, S., et al. 2020. Anti-stress, glial- and neuro-differentiation potential of resveratrol: characterization by cellular, biochemical and imaging assays. Nutrients 12: 671.
- Kaul, A., et al. 2021. Molecular insights into the antistress potentials of Brazilian green propolis extract and its constituent artepillin C. Molecules 27: 80.
- Gavid, M., et al. 2023. Technique of flat-mount immunostaining for mapping the olfactory epithelium and counting the olfactory sensory neurons. PLoS ONE 18: e0280497.

RESEARCH USE

For research use only, not for use in diagnostic procedures.