

Endoglin siRNA (m): sc-35303

BACKGROUND

Hereditary hemorrhagic telangiectasia (HHT) is an autosomal dominant disorder characterized by vascular abnormalities such as dilated vessels, hemorrhages, liver and lung congestion, and brain or heart ischemia. Mutations in two genes, Endoglin (also designated CD105) and ALK-1 (Activin receptor-like kinase-1, also designated TGF β superfamily RI), are responsible for HHT. Endoglin is mutated in HHT1 and ALK-1 is mutated in HHT2, both of which are thought to be caused by haploinsufficiency. Endoglin and ALK-1 are type III and type I members of the TGF β receptor superfamily, respectively, that are expressed on vascular endothelial cells. Endoglin can only bind ligands of the TGF β superfamily via association with the respective ligand binding receptors for TGF β 1, TGF β 3, Activin A, BMP-2 and BMP-7. The human ALK-1 gene encodes two protein species which exist as a result of either glycosylation or alternative splicing events. ALK-1 preferentially binds TGF β 1 and is expressed in bone marrow stromal cells, lung, brain, kidney and spleen.

REFERENCES

1. Wu, X., et al. 1995. Cloning and characterization of the murine activin receptor like kinase-1 (ALK-1) homolog. *Biochem. Biophys. Res. Commun.* 216: 78-83.
2. Altomonte, M., et al. 1996. Expression and structural features of Endoglin (CD105), a transforming growth factor β 1 and β 3 binding protein, in human melanoma. *Br. J. Cancer* 74: 1586-1591.
3. Gallione, C.J., et al. 1998. Mutation and expression analysis of the Endoglin gene in hereditary hemorrhagic telangiectasia reveals null alleles. *Hum. Mutat.* 11: 286-294.

CHROMOSOMAL LOCATION

Genetic locus: Eng (mouse) mapping to 2 B.

PRODUCT

Endoglin siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Endoglin shRNA Plasmid (m): sc-35303-SH and Endoglin shRNA (m) Lentiviral Particles: sc-35303-V as alternate gene silencing products.

For independent verification of Endoglin (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-35303A, sc-35303B and sc-35303C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Endoglin siRNA (m) is recommended for the inhibition of Endoglin expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Endoglin (P3D1): sc-18838 is recommended as a control antibody for monitoring of Endoglin gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Endoglin gene expression knockdown using RT-PCR Primer: Endoglin (m)-PR: sc-35303-PR (20 μ l, 580 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

1. Lee, S.B., et al. 2016. Identification of a distinct subpopulation of fibroblasts from murine dermis: CD73⁺ CD105⁺ as potential marker of dermal fibroblasts subset with multipotency. *Cell Biol. Int.* 40: 1008-1016.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.