

GADD 45 α siRNA (h): sc-35440

BACKGROUND

It is well established that cell cycle progression is subject to arrest at G₁ and G₂ checkpoints in response to DNA damage, presumably to allow time for DNA repair prior to entry into S and M phase, respectively. The p53 tumor suppressor is required for one such G₁ checkpoint and functions to upregulate expression of GADD 45 and p21. p21 functions to inhibit the kinase activity of multiple Cdk complexes, which may account for its suppression of cell growth. GADD 45 binds both Cdks and PCNA, a protein involved in DNA replication and repair. GADD 45 has been shown to stimulate DNA excision repair *in vitro* and to inhibit entry of cells into S phase. Thus, it has been suggested that GADD 45 may serve as a link between p53-dependent cell cycle checkpoint and DNA repair.

CHROMOSOMAL LOCATION

Genetic locus: GADD45A (human) mapping to 1p31.3.

PRODUCT

GADD 45 α siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see GADD 45 α shRNA Plasmid (h): sc-35440-SH and GADD 45 α shRNA (h) Lentiviral Particles: sc-35440-V as alternate gene silencing products.

For independent verification of GADD 45 α (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-35440A, sc-35440B and sc-35440C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

GADD 45 α siRNA (h) is recommended for the inhibition of GADD 45 α expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

GADD 45 α (C-4): sc-6850 is recommended as a control antibody for monitoring of GADD 45 α gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor GADD 45 α gene expression knockdown using RT-PCR Primer: GADD 45 α (h)-PR: sc-35440-PR (20 μ l, 498 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

1. Meyer, N.J., et al. 2009. Gadd45a is a novel candidate gene in inflammatory lung injury via influences on Akt signaling. *FASEB J.* 23: 1325-1337.
2. Asuthkar, S., et al. 2011. Gadd45a sensitizes medulloblastoma cells to irradiation and suppresses MMP-9-mediated EMT. *Neuro Oncol.* 13: 1059-1073.
3. Godel, M., et al. 2020. Small nucleolar RNAs determine resistance to doxorubicin in human osteosarcoma. *Int. J. Mol. Sci.* 21: 4500.
4. Jung, S.N., et al. 2021. EGR1/GADD 45 α activation by ROS of non-thermal plasma mediates cell death in thyroid carcinoma. *Cancers* 13: 351.
5. Jang, H.J., et al. 2021. Chelidonine induces apoptosis via GADD45A-p53 regulation in human pancreatic cancer cells. *Integr. Cancer Ther.* 20: 15347354211006191.
6. Lin, H.Y., et al. 2021. Epigenetic therapy combination of UNC0638 and CI-994 suppresses breast cancer via epigenetic remodeling of BIRC5 and GADD45A. *Biomed. Pharmacother.* 145: 112431.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.