

# HSP 90 $\beta$ siRNA (m): sc-35607

## BACKGROUND

Heat shock protein (HSP) molecular chaperones are environmental stress-inducible gene products. The human HSP 90 family includes 17 genes that fall into four classes: HSP90AA, HSP90AB, HSP90B and TRAP. HSP 90 family members guide the normal folding, intracellular disposition and proteolytic turnover of many key regulators of cell growth, differentiation and survival. HSP 90 $\alpha$ , also designated HSP 90A, HSP 86 and LPS-associated protein 2 (LAP2), is a cytosolic enhancer of inducible nitric-oxide synthase (iNOS), with chaperone activity that is important for the transcriptional activity of p53. HSP 90 $\beta$ , also designated HSP 90B, HSP 84 and HSPC2, is a cytosolic protein that participates in signaling pathways with PKC  $\epsilon$  to protect cells from external damage, particularly in heat shock-mediated events. GRP 94, also known as tumor rejection antigen 1 (TRA1), ECGP and GP96, localizes to the ER, is highly expressed in BGC-823 human gastric carcinoma cells and is upregulated in human endothelial cells in response to hypoxia by HIF-1. TRAP1 (TNF receptor-associated protein 1), also designated HSP 75) is a mitochondrial matrix component that plays a role in the induction of apoptosis in response to reactive oxygen species.

## REFERENCES

1. Wu, J.M., et al. 2003. PKC  $\epsilon$  is a unique regulator for HSP 90 $\beta$  gene in heat shock response. *J. Biol. Chem* 278: 51143-51149.
2. Whitesell, L., et al. 2005. HSP 90 and the chaperoning of cancer. *Nat. Rev. Cancer* 5: 761-772.
3. Cowen, L.E., et al. 2005. HSP 90 potentiates the rapid evolution of new traits: drug resistance in diverse fungi. *Science* 309: 2185-2189.

## CHROMOSOMAL LOCATION

Genetic locus: Hsp90ab1 (mouse) mapping to 17 B3.

## PRODUCT

HSP 90 $\beta$  siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see HSP 90 $\beta$  shRNA Plasmid (m): sc-35607-SH and HSP 90 $\beta$  shRNA (m) Lentiviral Particles: sc-35607-V as alternate gene silencing products.

For independent verification of HSP 90 $\beta$  (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-35607A, sc-35607B and sc-35607C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

HSP 90 $\beta$  siRNA (m) is recommended for the inhibition of HSP 90 $\beta$  expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

HSP 90 $\alpha/\beta$  (F-8): sc-13119 is recommended as a control antibody for monitoring of HSP 90 $\beta$  gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor HSP 90 $\beta$  gene expression knockdown using RT-PCR Primer: HSP 90 $\beta$  (m)-PR: sc-35607-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## SELECT PRODUCT CITATIONS

1. Zhang, M., et al. 2018. Jujuboside A promotes A $\beta$  clearance and ameliorates cognitive deficiency in Alzheimer's disease through activating Axl/HSP90/PPAR $\gamma$  pathway. *Theranostics* 8: 4262-4278.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.