

karyopherin α 2 siRNA (h): sc-35741

BACKGROUND

Protein transport across the nucleus is a selective, multi-step process involving several cytoplasmic factors. Proteins must be recognized as import substrates, dock at the nuclear pore complex and translocate across the nuclear envelope in an ATP-dependent fashion. Two cytosolic factors centrally involved in the recognition and docking process are the karyopherin α 1 and karyopherin β 1 subunits. Karyopherin α 1 functions in the recognition and targeting of substrates destined for nuclear import, while karyopherin β 1 serves as an adapter, tethering the karyopherin α 1/substrate complex to docking proteins on the nuclear envelope termed nucleoporins. Karyopherin α 2 has been shown to complex with Epstein-Barr virus nuclear antigen 1 (EBNA1). Certain RNA-binding proteins are imported to the nucleus by karyopherin β 2, and karyopherin β 3 appears to be involved in the import of some ribosomal proteins.

REFERENCES

1. Moroianu, J., et al. 1995. Previously identified protein of uncertain function is karyopherin α and together with karyopherin β docks import substrate at nuclear pore complexes. *Proc. Natl. Acad. Sci. USA* 92: 2008-2011.
2. Moroianu, J., et al. 1995. Protein export from the nucleus requires the GTPase Ran and GTP hydrolysis. *Proc. Natl. Acad. Sci. USA* 92: 4318-4322.

CHROMOSOMAL LOCATION

Genetic locus: KPNA2 (human) mapping to 17q24.2.

PRODUCT

karyopherin α 2 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see karyopherin α 2 shRNA Plasmid (h): sc-35741-SH and karyopherin α 2 shRNA (h) Lentiviral Particles: sc-35741-V as alternate gene silencing products.

For independent verification of karyopherin α 2 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-35741A, sc-35741B and sc-35741C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

karyopherin α 2 siRNA (h) is recommended for the inhibition of karyopherin α 2 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

karyopherin α 2 (B-9): sc-55538 is recommended as a control antibody for monitoring of karyopherin α 2 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor karyopherin α 2 gene expression knockdown using RT-PCR Primer: karyopherin α 2 (h)-PR: sc-35741-PR (20 μ l, 449 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

1. van der Watt, P.J., et al. 2009. The karyopherin proteins, Crm1 and karyopherin β 1, are overexpressed in cervical cancer and are critical for cancer cell survival and proliferation. *Int. J. Cancer* 124: 1829-1840.
2. Cheng, F., et al. 2010. Regulation of cell proliferation by the opioid growth factor receptor is dependent on karyopherin β and Ran for nucleocytoplasmic trafficking. *Exp. Biol. Med.* 235: 1093-1101.
3. Badding, M.A., et al. 2013. Proteomic and functional analyses of protein-DNA complexes during gene transfer. *Mol. Ther.* 21: 775-785.
4. Lin, J., et al. 2015. MiR-26b/KPNA2 axis inhibits epithelial ovarian carcinoma proliferation and metastasis through downregulating OCT4. *Oncotarget* 6: 23793-23806.
5. Lin, K.C., et al. 2018. Graphene oxide sensitizes cancer cells to chemotherapeutics by inducing early autophagy events, promoting nuclear trafficking and necrosis. *Theranostics* 8: 2477-2487.
6. Qin, L., et al. 2021. CCM3 loss-induced lymphatic defect is mediated by the augmented VEGFR3-ERK1/2 signaling. *Arterioscler. Thromb. Vasc. Biol.* 41: 2943-2960.
7. Jiang, L., et al. 2022. Decreased expression of karyopherin α 1 is related to the malignant degree of cervical cancer and is critical for the proliferation of Hela cells. *Pathol. Oncol. Res.* 28: 1610518.

RESEARCH USE

For research use only, not for use in diagnostic procedures.