

Laminin β -2 siRNA (m): sc-35785

BACKGROUND

The laminins comprise a growing family of disulfide-linked heterotrimers consisting of three genetically distinct polypeptide chains, designated α , β and γ . A major component of the basal lamina, laminins play a crucial role in providing a scaffolding upon which tissues are assembled and which serves as a physical barrier separating specialized tissues. During embryogenesis and early development, cells migrate along basement membranes, which are required for the polarization of cells. The β -2 Laminin chain plays a role in melanoma spread, promoting tumor migration along the abluminal surface of a vessel, a phenomenon which has been termed extra-vascular migratory metastasis.

REFERENCES

1. Yurchenco, P.D., et al. 1994. Basal lamina assembly. *Curr. Opin. Cell Biol.* 6: 674-681.
2. Engvall, E. 1995. Structure and function of basement membranes. *Int. J. Dev. Biol.* 39: 781-787.
3. Aumailley, M., et al. 1996. Laminins: a family of diverse multifunctional molecules of basement membranes. *J. Invest. Dermatol.* 106: 209-214.
4. Nomizu, M., et al. 1996. Mechanism of Laminin chain assembly into a triple-stranded coiled-coil structure. *Biochemistry* 35: 2885-2893.
5. Ziober, B.L., et al. 1996. Laminin-binding integrins in tumor progression and metastasis. *Semin. Cancer Biol.* 7: 119-128.
6. Ancsin, J.B., et al. 1996. Laminin interactions important for basement membrane assembly are promoted by zinc and implicate Laminin zinc finger-like sequences. *J. Biol. Chem.* 271: 6845-6851.
7. Durkin, M.E., et al. 1996. Structural organization of the human and mouse Laminin β -2 chain genes, and alternative splicing at the 5' end of the human transcript. *J. Biol. Chem.* 271: 13407-13416.
8. Vogel, W., et al. 1999. Expression of Laminin β -2 chain in normal human bone marrow. *Blood* 94: 1143-1145.
9. Libby, R.T., et al. 1999. Disruption of Laminin β -2 chain production causes alterations in morphology and function in the CNS. *J. Neurosci.* 19: 9399-9411.

CHROMOSOMAL LOCATION

Genetic locus: Lamb2 (mouse) mapping to 9 F2.

PRODUCT

Laminin β -2 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Laminin β -2 shRNA Plasmid (m): sc-35785-SH and Laminin β -2 shRNA (m) Lentiviral Particles: sc-35785-V as alternate gene silencing products.

For independent verification of Laminin β -2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-35785A, sc-35785B and sc-35785C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Laminin β -2 siRNA (m) is recommended for the inhibition of Laminin β -2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Laminin β -2 (C4): sc-59980 is recommended as a control antibody for monitoring of Laminin β -2 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Laminin β -2 gene expression knockdown using RT-PCR Primer: Laminin β -2 (m)-PR: sc-35785-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.