

MafB siRNA (m): sc-35840

BACKGROUND

Members of the Maf family of basic region/leucine zipper (bZIP) transcription factors affect transcription in either a positive or negative fashion, depending on their particular protein partner and the context of the target promoter. c-Maf (Maf-2) and the closely related family members Neural retina leucine zipper (Nrl), L-Maf, and Krm1/MafB (Maf-1) all bind to T-MARE sites and have been implicated in a wide variety of developmental and physiologic roles. The three small Maf family proteins (MafF, MafG, and MafK) are components of NF-E2 that function as heterodimers with the large tissue-restricted subunit of NF-E2 called p45, and they are implicated in the transcriptional regulation of many erythroid-specific genes. MafB is expressed in a wide variety of tissues and encodes a protein containing a typical bZip motif in its carboxy-terminal region. As a transcriptional activator, MafB plays a pivotal role in regulating lineage-specific gene expression during hematopoiesis by repressing Ets-1-mediated transcription of key erythroid-specific genes in myeloid cells. c-Maf interacts with the c-Myb DNA binding domain and forms Myb-Maf complexes, which in turn mediate the cooperative interactions between c-Myb and Ets-1 during early myeloid cell differentiation.

REFERENCES

1. Kerppola, T.K., et al. 1994. A conserved region adjacent to the basic domain is required for recognition of an extended DNA binding site by Maf/Nrl family proteins. *Oncogene* 9: 3149-3158.
2. Igarashi, K., et al. 1995. Conditional expression of the ubiquitous transcription factor MafK induces erythroleukemia cell differentiation. *Proc. Natl. Acad. Sci. USA* 92: 7445-7449.

CHROMOSOMAL LOCATION

Genetic locus: MafB (mouse) mapping to 2 H2.

PRODUCT

MafB siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see MafB shRNA Plasmid (m): sc-35840-SH and MafB shRNA (m) Lentiviral Particles: sc-35840-V as alternate gene silencing products.

For independent verification of MafB (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-35840A, sc-35840B and sc-35840C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

MafB siRNA (m) is recommended for the inhibition of MafB expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

MafB (B-11): sc-376387 is recommended as a control antibody for monitoring of MafB gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor MafB gene expression knockdown using RT-PCR Primer: MafB (m)-PR: sc-35840-PR (20 μ l, 561 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

1. Johnson, N.B., et al. 2016. Perturbed MafB/GATA1 axis after burn trauma bares the potential mechanism for immune suppression and anemia of critical illness. *J. Leukoc. Biol.* 100: 725-736.
2. Meng, Y., et al. 2020. MicroRNA-148a facilitates inflammatory dendritic cell differentiation and autoimmunity by targeting MAFB. *JCI Insight* 5: e133721.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.