



MC5-R siRNA (m): sc-35876

BACKGROUND

The melanocortin family comprises the α -, β - and γ -melanocyte stimulating hormones (MSH) and adrenocorticotrophin. The receptors for these hormones are seven-transmembrane G protein-coupled proteins that activate adenyl cyclase. Five melanocortin receptors have been cloned and shown to exhibit different affinities and patterns of expression. MC1-R (MSH-R) is expressed in melanocytes and corticoadrenal tissue. MC2-R is the ACTH receptor and is expressed primarily in the adrenal cortex. MC3-R has been found in specific regions of the brain and is also expressed in placenta and gut. MC4-R is expressed primarily in brain, while MC5-R is expressed at low levels in most tissues.

REFERENCES

1. Mountjoy, K.G., et al. 1992. The cloning of a family of genes that encode the melanocortin receptors. *Science* 257: 1248-1251.
2. Chhajlani, V., et al. 1992. Molecular cloning and expression of the human melanocyte stimulating hormone receptor cDNA. *FEBS Lett.* 309: 417-420.
3. Gantz, I., et al. 1993. Molecular cloning of a novel melanocortin receptor. *J. Biol. Chem.* 268: 8246-8250.
4. Roselli-Reh fuss, L., et al. 1993. Identification of a receptor for γ melanotropin and other proopiomelanocortin peptides in the hypothalamus and limbic system. *Proc. Natl. Acad. Sci. USA* 90: 8856-8860.
5. Gantz, I., et al. 1993. Molecular cloning, expression and gene localization of a fourth melanocortin receptor. *J. Biol. Chem.* 268: 15174-15179.
6. Chhajlani, V., et al. 1993. Molecular cloning of a novel human melanocortin receptor. *Biochem. Biophys. Res. Commun.* 195: 866-873.
7. Mountjoy, K.G., et al. 1994. Localization of the melanocortin-4 receptor (MC4-R) in neuroendocrine and autonomic control circuits in the brain. *Mol. Endocrinol.* 8: 1298-1308.
8. Labbe, O., et al. 1994. Molecular cloning of a mouse melanocortin 5 receptor gene widely expressed in peripheral tissues. *Biochemistry* 33: 4543-4549.

CHROMOSOMAL LOCATION

Genetic locus: Mc5r (mouse) mapping to 18 E2.

PRODUCT

MC5-R siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see MC5-R shRNA Plasmid (m): sc-35876-SH and MC5-R shRNA (m) Lentiviral Particles: sc-35876-V as alternate gene silencing products.

For independent verification of MC5-R (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-35876A, sc-35876B and sc-35876C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

MC5-R siRNA (m) is recommended for the inhibition of MC5-R expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor MC5-R gene expression knockdown using RT-PCR Primer: MC5-R (m)-PR: sc-35876-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

1. Shimizu, K., et al. 2015. Histamine released from epidermal keratinocytes plays a role in α -melanocyte-stimulating hormone-induced itching in mice. *Am. J. Pathol.* 185: 3003-3010.

PRESEARCH USE

For research use only, not for use in diagnostic procedures.

ROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.