

# INTERVET INC., MERCK ANIMAL HEALTH

**556 MORRIS AVE., SUMMIT, NJ, 07901**

Telephone: 862-245-4321  
Order Desk: 800-648-2118  
Fax: 862-245-4935  
Customer Service: 800-521-5767  
Website: [www.intervet.com](http://www.intervet.com)



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the USA product label or package insert.

## **NOBIVAC® PUPPY-DPv**

***Intervet/Schering-Plough Animal Health***

**Canine Distemper-Parvovirus Vaccine**

**MODIFIED LIVE VIRUS**

For veterinary use in dogs only

**IN THE ABSENCE OF A VETERINARIAN-CLIENT-PATIENT RELATIONSHIP, U.S. FEDERAL REGULATIONS PROHIBIT THE RELABELING, REPACKAGING, RESALE OR REDISTRIBUTION OF THE INDIVIDUAL CONTENTS OF THIS PACKAGE.**

THIMEROSAL FREE

### **PRODUCT DESCRIPTION:**

Nobivac® Puppy-DPv is a modified live virus vaccine containing attenuated strains of canine distemper virus (CDV) and parvovirus (CPV). This vaccine is for vaccination of healthy dogs as an aid in the prevention of disease caused by canine distemper virus and canine parvovirus.

### **DOSAGE AND ADMINISTRATION:**

Aseptically reconstitute the desiccated vaccine with the sterile diluent provided and administer 1mL by the subcutaneous route or intramuscular route. Initial vaccination of healthy dogs may be given as early as 4 weeks of age, with booster injections administered every 3-4 weeks until the dog is 12 weeks of age. Dogs over 12 weeks of age should initially receive 2 doses 3-4 weeks apart. Annual revaccination with one dose is recommended.

Following the advice of the American Veterinary Medical Association on Vaccination Principles, an appropriate revaccination program for individual animals should be made based upon veterinarian-client-patient relationships.

### **PRECAUTIONS:**

- Store at 2° - 7°C (35°-45°F).
- Do not freeze.
- Shake well before use.
- Use contents promptly once reconstituted.
- Use new, non-chemically sterilized needles and syringes.
- Burn vaccine container and all unused contents.
- Contains gentamicin and amphotericin B as preservatives.
- Vaccinate only healthy, non-parasitized dogs.
- Do not vaccinate pregnant bitches.
- The use of a biological product may produce anaphylaxis and/or other inflammatory immune-mediated hypersensitivity reactions. **Antidote:** Epinephrine, corticosteroids, and antihistamines may all be indicated depending on the nature and severity of the reaction.

Intervet Inc., Omaha, NE 68103 USA

U.S. Veterinary License No. 165A

1 800 224-5318 (USA)

1 888 306-0069 (Canada)

© 2010 Intervet Inc.

All rights reserved.

CPV STRAIN 154®

U.S. Patent No.

4,810,494

5,316,764

	Code	
25x1 DOSE	053260	110740-011 AL302

**NAC No.:** 10473531