



# Orexin R-2 siRNA (m): sc-36134

## BACKGROUND

The hypothalamus is essential for maintaining homeostasis by integrating the vertebrate endocrine and nervous systems, thereby controlling temperature, thirst and hunger. Orexin-A and Orexin-B (also designated hypocretins) are hypothalamic neuropeptides that are derived from a single precursor, prepro-orexin, by proteolytic processing. These peptides bind to and activate two closely related, G protein-coupled receptors, designated Orexin receptor-1 (Orexin R-1) and Orexin receptor-2 (Orexin R-2). Orexin-A protein and prepro-orexin mRNA are localized to neurons within the lateral section of the hypothalamus, designated the "feeding center". Prepro-orexin mRNA is up-regulated during fasting, suggesting that orexins may play a role in the central feedback mechanism that regulates feeding behavior. Orexin has been shown to increase the release of GABA and glutamate from axons, a response seen as a result of most synaptic activities in the hypothalamic region.

## REFERENCES

1. Campbell, N. 1990. Biology 2nd ed. Redwood City: The Benjamin/Cummings Publishing Company, Inc., 916-921.
2. Sakurai, T., et al. 1998. Orexins and orexin receptors: a family of hypothalamic neuropeptides and G protein-coupled receptors that regulate feeding behavior. *Cell* 92: 573-585.
3. de Lecea, L., et al. 1998. The hypocretins: hypothalamus-specific peptides with neuroexcitatory activity. *Proc. Natl. Acad. Sci. USA* 95: 322-327.
4. Wolf, G. 1998. Orexins: a newly discovered family of hypothalamic regulators of food intake. *Nutr. Rev.* 56: 172-173.
5. van der Pol, A., et al. 1998. Presynaptic and postsynaptic actions and modulation of neuroendocrine neurons by a new hypothalamic peptide, hypocretin/orexin. *J. Neurosci.* 18: 7962-7971.
6. Yamanaka, A., et al. 2002. Orexins activate histaminergic neurons via the orexin 2 receptor. *Biochem. Biophys. Res. Commun.* 290: 1237-1245.
7. Harris, D.M., et al. 2002. Stimulation of amylase release by Orexin is mediated by Orexin 2 receptor in AR42J cells. *Pancreas* 25: 405-410.
8. Lang, M., et al. 2004. Structure-activity studies of orexin A and orexin B at the human Orexin 1 and Orexin 2 receptors led to Orexin 2 receptor selective and Orexin 1 receptor preferring ligands. *J. Med. Chem.* 47: 1153-1160.
9. Beiras-Fernandez, A., et al. 2004. Merkel cells, a new localization of prepro-Orexin and Orexin receptors. *J. Anat.* 204: 117-122.

## CHROMOSOMAL LOCATION

Genetic locus: Hcrtr2 (mouse) mapping to 9 D.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## PRODUCT

Orexin R-2 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Orexin R-2 shRNA Plasmid (m): sc-36134-SH and Orexin R-2 shRNA (m) Lentiviral Particles: sc-36134-V as alternate gene silencing products.

For independent verification of Orexin R-2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-36134A, sc-36134B and sc-36134C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

Orexin R-2 siRNA (m) is recommended for the inhibition of Orexin R-2 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Orexin R-2 gene expression knockdown using RT-PCR Primer: Orexin R-2 (m)-PR: sc-36134-PR (20  $\mu$ l, 587 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.